

DETECTING METHOD AND DETECTION SUBSTRATE  
FOR USE THEREIN

BACKGROUND OF THE INVENTION

5 Field of the Invention

The present invention is directed to examining multiple specimens at a time for multiple items, and provides a method in which matrix substrates with biological samples having different properties and origins bound thereto are prepared, and on each matrix region, oligonucleotides having different sequences, proteins or drugs are spotted in an array, whereby multiple specimens are examined at a time for multiple items.

10 The present invention also relates to a method in which by using an oligonucleotide having a known base sequence as a detection probe to detect whether a complex is formed by intermolecular bond with this oligonucleotide, detection is made whether or not components having a capability of bonding to the above described detection probe are contained, and a detection substrate having the oligonucleotide as a detection probe fixed on its surface, which is used exclusively for this detecting method.

25 Related Background Art

In identification of partial sequences included in the base sequence of a nucleic acid molecule, detection

of a target nucleic acid contained in a sample  
originated from an organism or identification of genus  
or species for various bacteria based on the  
characteristics of the gene DNA of the bacteria, a  
5 procedure may be used in which two or more probe DNAs  
having known base sequences are used to detect whether  
or not the nucleic acid molecule is a nucleic acid  
molecule specifically binding to each probe DNA, namely  
making hybridization with each probe DNA. As an  
10 effective approach to performing speedily and  
accurately examination of the two or more probe DNAs by  
the hybridization method, a procedure is proposed in  
which a probe array of two or more probe DNAs arranged  
regularly on a solid phase is used to detect at a time  
15 whether or not the nucleic acid molecule is a nucleic  
acid molecule specifically binding to each probe DNA.

Among common methods for producing such probe  
arrays, as described in European Patent No. 373203 (EP  
0373203 B1) for example, methods are known in which  
20 predetermined nucleic acid probes are synthesized in an  
array form on a solid phase, and methods in which a  
plurality of nucleic acid probes synthesized in advance  
is supplied in an array form on the solid phase.

Prior technical documents disclosing the former  
25 methods include, for example, U.S. Patent No.  
5,405,783. Also, as one example of the latter methods,  
a method in which cDNAs are arranged in an array form

on a solid phase using micropipetting is disclosed in, for example, U.S. Patent No. 5,601,980 and "Science", Vol. 270, pp. 467, (1995).

5       The probe array that is prepared with these  
methods may be an array such that nucleic acid probes  
are arranged on a solid phase at a high density of  
10000 or more probes per square inch. Hybridization  
reaction with multiple probes are carried out at a time  
by dipping this high-density probe array into a  
10   specimen solution, and in so doing, the base sequence  
of genes is analyzed based on the base sequence of  
nucleic acids making hybridization. This method has an  
advantage that probes are arranged in a high density on  
a substrate of small area, thereby making it possible  
15   to conduct multiple-item examination at a time with a  
small amount of samples to reduce burden associated  
with sampling from the subject.

As a method of preparing the high-density probe  
array for the above described application on the  
20   substrate by the DNA synthesis process, a method in  
which a photolithography technology is applied is  
disclosed in the aforesaid U.S. Patent No. 5,405,783,  
but highly advanced equipment is required for  
implementing this method, and the method is not easy  
25   enough for anyone to use.

Also, in the case where the number of specimens is  
large but the number of required examination items is

not so large, the integration degree of DNA probes on the probe array corresponding to the number of examination items does not need to be very high.

Rather, there may be cases where it is necessary to  
5 prepare a large number of probe arrays with a small number of desired DNA probes fixed, using a simpler method.

Actually, in the field of clinical examination, there are not necessarily many cases where examinations  
10 for more than 10000 items are required. For example, in the case of group health examination and the like, there may be cases where it is more important to examine a large number of specimens with a limited number of items. For examining a large number of  
15 specimens in this way, a system is required such that presence of diseases can be speedily examined through comparison with standard samples with respect to each specimen.

In addition, the amount of DNA specimen is  
20 generally small as compared to that of oligonucleotide capable of being synthesized and used in the probe. For using it in a normal form in which the probe array substrate is dipped into the specimen solution for hybridization reaction, the amount of specimen DNA  
25 allowing the substrate to be dipped sufficiently is required. Therefore, the size of the DNA probe array substrate is limited depending on the amount of

specimen DNA, and thus the array needs to be highly dense. Alternatively, as a result of diluting the specimen solution to ensure its volume for the size of the probe array substrate, the concentration of DNA in the specimen solution is reduced, and a procedure is adopted of prolonging reaction time to make compensation for the reduced concentration.

Also, since the amount of sampled specimens is limited inherently because the specimen is an extract from tissues, and it is subjected to pre-processing for making a specimen solution for use in hybridization reaction, specifically extraction of nucleic acid, single-strand formation thereof, and process for labeling, the amount of finally obtained samples is very small. In order to make compensation for that, the sample is subjected to processing for amplification of the amount of DNA such as amplification processing by PCR reaction before it is used for examination and studies. However, there exists a disadvantage that because primers separately prepared are required for carrying out a PCR reaction, such processing can be applied only to specific genes of which primer sequence is known. In addition, there exist sequences that can easily be amplified and sequences that can hardly be amplified in the process of PCR reaction, and thus the efficiency of reaction (rate of amplification) is not uniform. For example, in the case where the content of

a specific mRNA in the total amount of extracted mRNA is examined to determine diseases or like based on the content, standard samples providing criteria should be always prepared to make correction on the above

5 described amplification rate.

Although the amount of the specimen solution required for hybridization reaction decreases as the size of the substrate is reduced, there is a limitation on downsizing of the substrate in association with  
10 handling. Specifically, it is possible in principle to enhance array density and reduce the number of probes to be placed on the array to downsize the substrate, but if an extremely small substrate is used, a dedicated handling apparatus is required in the process  
15 of processing such as hybridization reaction and detection thereafter, which cannot be practical.

Also, for examining cDNA for mRNA that is transcribed with reflection of the process of development of a certain organism, cDNA for mRNA that  
20 is transcribed with reflection of each phase in the process of culturing a certain cell, cDNA for mRNA that is transcribed by interaction with drugs, and so on, a DNA array with multiple types of test samples arranged is used. Examples of arraying this test sample are  
25 described, for example, in the above described "Science", Vol. 270, pp 460, (1995). In this case, test samples arrayed on the substrate are dipped using

as a probe solution the labeled DNA of known sequence that is derived from genes having a specific function, whereby hybridization reaction is carried out.

5 If a plurality of items is to be examined at a time using this methodology, DNA probes labeled with different types of fluorescent reagents (fluorochromes) should be prepared depending on the number of items. When detection is made, those different types of fluorescent reagents (fluorochromes) must be detected  
10 as distinguished from one another, and therefore their wavelengths and the like should be different as a matter of course. Of course, detection filters corresponding to respective fluorescent reagents (fluorochromes) are also needed for a detector.

15 This need for simultaneous examination of multiple items for multiple specimens is not characteristic exclusively of hybridization reaction among genes (DNA).

For example, it is also important to examine  
20 multiple items with a small amount of samples as to interaction between genes and other substances such as interaction between genes and proteins (DNA binding proteins) and screening of chemicals that are bound to genes. Detection of former DNA binding proteins is  
25 used to elucidate the control mechanism of gene expression by proteins such as transcription accelerators, but in the present situation, methods in

which DNA fragments are bound to proteins, and thereafter complexes are analyzed by gel electrophoresis are adopted. In this method, the number of specimens that can be analyzed at a time is limited due to usage of gel electrophoresis, and considerable time is required for analysis.

For the field of development of drugs, there may be cases where examination of interaction between genes and administered drugs constitutes an important item in progress of research, but it takes relatively much time and efforts to obtain chemically synthesized products for use in drugs to be researched, and it can be considered that reduction in the amount of drugs to be used in screening results in significant improvement in efficient of their research.

As introduced above, there are cases where when a complex is formed using interaction between two substances such as hybridization between DNAs, formation of a complex of DNA and a protein, and interaction of a drug compound with gene DNA, or the presence or absence of interaction causing a complex to be formed is examined, the amount of samples of one of those two substances is limited, and the limited amount of samples should be used to conduct a series of desired examinations across multiple types as to the presence or absence of formed complexes. That is, development of an examination method in which



consumption of samples required for individual examinations can be reduced to carry out examination across multiple types more efficiently within a limited amount of samples is desired.

5

#### SUMMARY OF THE INVENTION

10 An object of the first invention is to provide a method of examining multiple specimens at a time for multiple items, for example a method in which matrix substrates with biological samples having different properties and origins bound thereto are prepared, and on each matrix region, oligonucleotides or proteins having different sequences and drugs are spotted in an array form, whereby multiple specimens are examined at  
15 a time for multiple items.

Another object of the invention is to provide a method in which multiple specimens can also be examined at a time for multiple items in a similar way for interaction between chemicals, especially drugs, and  
20 cDNA, binding of proteins to cDNA and the like.

An object of the second invention is to provide a new method in which oligonucleotide of which base sequence is known and which can be obtained relatively easily is used as a detection probe, and when for a  
25 limited amount of sampled specimens, the presence or absence of a bonding capability to the above described oligonucleotide as a detection probe or the degree of

the bonding capability is examined by the presence or absence of complexes formed between those two substances, or efficiency thereof is evaluated, consumption of specimens required for evaluation for  
5 each type of oligonucleotide as a detection probe can be reduced. In addition, the invention also has an object to provide a detection substrate with the above described oligonucleotide being fixed as a detection  
10 probe in a predetermined region of its surface, which is used exclusively for the method, and provide a method of preparing the detection substrate.

The examination method of the first invention capable of achieving the above described objects is a method in which a reactivity between a first sample and  
15 a plurality of second samples having different properties from one another is examined at a time,

characterized in that in a defined region on a substrate with the first sample bound on the entire surface in advance, the second samples are placed  
20 independently of one another as spots having a smaller size than the above described defined region, and then the reactivity between the above described first sample and each of the second samples is tested.

The matrix of biological samples related to the  
25 invention that is usefully used for the above examination method is characterized in that two or more types of biological samples of different origins exist

in respective matrix regions separated on the substrate.

According to the invention, a substrate with biological samples having different properties and  
5 origins (e.g. nucleic acids and proteins) bound in a matrix form in advance can be provided.

There is also provided a method in which DNA probes like oligonucleotides, cDNAs, proteins or chemicals are spotted in an array form on the above  
10 described substrate with biological samples having different properties and origins placed in a matrix form to carry out reaction, and the presence or absence of another sample bound to a certain biological sample, the degree of the bonding, and the presence or absence  
15 of interaction is examined for multiple items at a time and speedily.

In this method, the area occupied by one specimen is very small because two or more types of specimens are placed on one substrate. Therefore, there is an  
20 advantage that the amount of required cDNA may be very small as compared to the case where hybridization reaction is carried out using a conventional DNA array with an enormously large number of DNA probes bound in an array form in advance. Also, there is neither  
25 limitation on the size of the DNA array substrate nor inconvenience for handling.

Also, by providing a method in which examination

can be carried out even with a small amount of samples,  
the method opens the door to areas in which examination  
could not be carried out because conventionally, a  
sufficient amount of samples cannot be obtained, for  
5 example a new examination area in which mRNA obtained  
from tissues is directly examined.

In addition, according to the invention, a method  
in which chemicals, proteins and nucleic acids can be  
examined at a time under the same reaction condition on  
10 the same substrate.

A method of detecting object components in test  
samples according to the second invention is a method  
in which using as a detection probe oligonucleotide of  
which base sequence is known, complexes formed between  
15 the above described oligonucleotide and the object  
components are detected to examine whether or not the  
object components having a capability of binding to the  
above described oligonucleotide are contained in the  
liquid test samples, or evaluate the degree of binding  
20 capability thereof,

characterized in that there is at least one type  
of the above described oligonucleotide used as a  
detection probe, of which base sequence is known,

there are at least two types of test samples to be  
25 examined, and

a detection substrate with the above described one  
or more types of oligonucleotide for detection probes

bound to predetermined sections respectively on a predetermined solid substrate is used,

the above described method comprising steps of:

5 spotting a plurality of predetermined amounts of sample solution for each spot so that a predetermined array shape is formed in the spotted position, for each of the above described two or more types of test samples, in each section with the oligonucleotide for detection probes bound in advance;

10 detecting the presence or absence of complexes formed between the above described oligonucleotide and the object component, for the above described plurality of spots for each test sample, respectively; and

15 determining whether or not the object component having a capability of binding to the above described oligonucleotide is contained, or the degree of the capability of binding, based on the result of the above described detection.

20 Also, the present invention provides a detection substrate that is exclusively used when the above described method of the invention is carried out. That is, the detection substrate of the present invention is a detection substrate with two or more oligonucleotides having known base sequences different from one another  
25 fixed on a solid substrate, characterized in that:

the above described plurality of oligonucleotides are bound and fixed in predetermined sections,

# BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows one example of an arrangement aspect of defined regions on a substrate in the present invention;

5        FIGS. 2A and 2B show one example of matrices in the present invention, wherein FIG. 2A is a plan view, and FIG. 2B is a 2B-2B sectional view thereof;

FIG. 3 is a schematic explanatory view of a specimen solution discharging method by bubble jet  
10       process that is an embodiment of the present invention;

FIG. 4 is a sectional view of a bubble jet head 105 taken in the 4-4 line in FIG. 3;

FIG. 5 shows a layout of 64 discharged DNA probes on each black matrix;

15       FIG. 6 shows one example of detection substrates of the present invention, illustrating schematically a situation in which sections in which oligonucleotides being detection probes are fixed are arranged in a matrix form, and a plurality of cDNAs are spotted in a  
20       two-dimensional array form onto each section as detection samples;

FIG. 7 illustrates schematically arrangements of respective probes in the detection substrate with 64 DNA probes bound to sections arranged in the form of a  
25       8 x 8 matrix, respectively;

FIG. 8 shows schematically a pattern of a spot array of total 64 x 64 in which 64 test samples are

respectively, so that one type of oligonucleotide exists in each section, and

5 a plurality of the above described sections in which oligonucleotides are fixed is placed in a matrix form on the surface of the above described solid substrate.

10 The method of preparing the detection substrate of the present invention is a method suitable for preparation of the above described detection substrate of the invention, and specifically is a method of preparing a detection substrate with two or more oligonucleotides having known base sequences different from one another fixed on a solid substrate, characterized in that:

15 for the above described solid substrate, a substrate with a plurality of sections separated in a matrix form in advance formed on the surface thereof is used,

20 the above described a plurality of oligonucleotides is supplied into predetermined sections in predetermined amounts using printing by ink jet process, respectively, so that one oligonucleotide is present in each section, and

25 the supplied oligonucleotides are fixed in the predetermined sections.

spotted in the form of a two-dimensional  $8 \times 8$  array on each section, for the detection substrate on which sections with probes fixed therein are arranged in the form of the  $8 \times 8$  matrix;

5        FIG. 9 shows schematically a result of spotting 64 test samples in the form of the two-dimensional  $8 \times 8$  array on each section for 64 probes fixed in sections arranged in the form of the  $8 \times 8$  matrix to carry out hybridization reaction; and

10        FIG. 10 shows an example of the structure of sections delimited by hydrophobic frame-structured walls provided on the detection substrate of the present invention, and arranged in the form of the  $8 \times 8$  matrix.

15

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

One embodiment of the present invention will be described below referring to FIG. 1. FIG. 1 shows a substrate surface with 64 defined regions formed thereon, wherein each region (matrix) measures 1 mm by 1 mm, and a space  $x$  between regions can be selected freely. For methods of preparing biological sample binding matrix substrates, for example, a method can be used in which the solution of a first sample (e.g. biological sample) is printed on the entire surface of defined regions on the substrate as a "solid print pattern" by coating and ink jet processes, or is

20

25



supplied by methods such as chemical synthesis on the substrate, and is bound in a matrix form on the substrate through adsorption to the substrate or chemical reaction between functional groups existing in the biological sample and functional groups existing on the substrate. Furthermore, the situation in which the first sample is bound on the entire surface of defined regions means a situation in which the first sample is bound across the entire surface such that when a second sample and samples thereafter are supplied in these defined regions, these reactions occur without being limited to the positions in the above described regions in which the samples are supplied. For example, the first sample may be fixed in layered form on the entire surface, or the masses of molecules constituting the first sample may be dispersed on the entire surface in high density with micro-spaces being kept among them.

The defined regions on this substrate may previously be provided on the substrate as a well constituted by sections separated in pattern form by walls of hydrophobic compounds.

Also, when using a substrate with nucleoside acid (cDNA) being a biological sample fixed thereon as the first sample, two or more probe DNAs possibly included in cDNA are contacted with cDNA on the substrate as the second sample and samples thereafter, and products of reaction with the above described probes are detected

on the above described solid phase to detect the presence or absence of probe DNA sequences in the above described cDNA, two or more probes are supplied in an array form as mutually independent spots in each matrix with various kinds of cDNA bound in the defined regions, thereby making it possible to perform simultaneous detection with two or more probes.

Also, on the nucleic acid (cDNA) matrix, two or more types of chemicals or proteins that are possibly bound to cDNA are contacted with the probe DNA on the substrate as mutually independent spots, thereby making it possible to perform multiple-item examination composed of these reactions at a time. Multiple-item screening of DNA binding proteins and DNA binding chemicals can be performed at a time by detecting presence of binding of chemicals or proteins to probes on the solid phase.

The present invention is characterized by supplying probe DNA, proteins and chemicals in a form of droplets of small amounts on the matrix on which biological samples such as cDNA are applied, wherein different types of samples are arranged in an array form, thereby making it possible to perform simultaneous multiple-item processing.

Combinations of the first sample fixed in advance on the substrate and the second sample and sample thereafter that are reacted with the first sample may

include the following combinations.

Specific examples of the matrix or the like formed of defined regions on the substrate for use in the present invention will be described below.

5           (Shapes of matrices with biological samples bound thereto)

10           The shapes of matrix patterns are not particularly limited, and may include any shapes, but shapes such as linear, squares and rectangular are preferable in that they can be treated irrespective of how specimens are supplied, in consideration of convenience at the time of supplying specimens on the created substrate. Of course, forms such as circles and ellipses will cause no problems.

15           Materials that are fixed to the substrate as a first sample may include unknown base sequences derived from organisms, cDNA libraries, mRNA libraries, sets of two or more DNA and RNA, known DNA and RNA synthesized or derived from organisms or sets thereof, chips of  
20           cloned oncogenes, protein fractions including at least one type of protein derived from organisms, proteins of single type, mixtures of known proteins of different types, and chemicals.

25           (Density of matrices with biological samples bound)

          The density of matrices is not particularly limited, but for a preferred form, the density of 400

per centimeter square is preferable. For density of 400/cm<sup>2</sup>, the size of one matrix is a 500 µm square in the case of square form. If samples to be arranged as spots on the array are arranged as spots with diameters of 100 µm, 25 spots are arranged in total with 5 spots high by 5 spots wide. Also, if the diameter of sample solution is 20 µm, the number of spots that can be arranged in a row is 25, and 625 spots can be arranged in total.

10           (Preparation of a substrate with biological samples bound thereto)

          Samples originated from organisms (biological samples) include nucleic acids and proteins. Nucleic acids include, for example, mRNA and cDNA, and methods for binding them on the substrate include a method in which nucleic acid extracted and purified in advance is applied to the substrate to fix the nucleic acid by adsorption and electrostatic bond, and a method in which the nucleic acid is fixed by providing covalent bond thorough chemical reaction with functional groups on the substrate using amino groups the nucleic acid has.

          The method using negative electric charges of DNA is a method in which nucleic acid is electrostatically bound to a solid carrier subjected to surface treatment with poly positive ions such as polylysine, polyethyleneimine and polyalkylamine, and then blocking

of excessive positive ions is carried out, which is generally used.

(Types of functional groups of solid phases and nucleic acids)

5 Combinations of functional groups that are used for fixation include, for example, a combination of epoxy groups (on solid phase) and amino groups (amino groups in nucleic acid probe terminals or base groups). Methods for introducing epoxy groups to the solid  
10 surface include, for example, a method in which polyglycidyl methacrylate having epoxy groups is applied to the solid surface composed of resin, and a method in which a silane coupling agent having epoxy groups is applied to the solid surface made of glass  
15 and is reacted with glass.

(Binding of proteins to the solid phase)

Methods of binding proteins to the substrate include methods using adsorption as in the case of nucleic acid and methods using electrostatic binding.  
20 Furthermore, methods of forming covalent bond include methods using SH groups of cysteine residues in addition to the above described methods using amino groups.


(Methods of fixation of proteins using thiol  
25 groups)

Methods using cysteine residues for fixation of proteins include, for example, methods using

combinations of maleimide groups and thiol groups (-SH). That is, treatment is done so that the solid surface has maleimide groups, whereby thiol groups of cysteine residues supplied to the solid surface can be  
5 reacted with maleimide groups of the solid surface to fix proteins.

For methods of introducing maleimide groups to the solid surface, a various kinds of methods may be used, and this can be achieved by, for example, reacting an  
10 aminosilane coupling agent with a glass substrate, and then reacting its amino groups with a reagent containing N-(6-maleimidocaproyloxy) succinimide) expressed by the following structural formula (EMCS reagent: manufactured by Dojin Co., Ltd.).

15



20

For another example, a reagent containing succinimidyl 4-(maleimidophenyl) butyrate can be used to react with amino groups preferably.

(DNA matrix structures composed of hydrophobic matrices)

25

For an additional form of fixation of biological samples, a method can be used in which a well composed of, for example, hydrophilic and hydrophobic matrices is formed on the solid surface, a structure to prevent

coupling among spots is provided in advance, and the DNA probe is supplied in the well to carry out coupling reaction.

(Materials of matrices/wells)

5           When probe solution is put on the separated matrix to carry out coupling reaction, it is preferable that portions constituting the well is hydrophilic, and portions corresponding to the wall surface of the well and the partition between the well and a neighboring  
10 well are composed of materials whose surfaces are less compatible with the probe solution. Due to such a treatment, the probe solution can be smoothly supplied to a desired well even if some positional deviation occurs when the probe solution is supplied to the well.

15           One example of matrices in this embodiment is shown in FIGS. 2A and 2B. FIG. 2A is a plan view, and FIG. 2B is a 2B-2B sectional view thereof. This matrix has a structure in which a matrix pattern 125 having a frame structure with formed recesses 127 (wells) placed  
20 in the form of a solid phase 103 is provided. The wells 127 separated from one another by the matrix 125 (height) are provided as through-holes (cut-off portions) in the matrix pattern, of which side is constituted by heights and of which bottom 129 has the  
25 exposed surface of the solid phase 103. The portion of the exposed surface of the solid phase 103 forms a surface that can be coupled to the probe, and the probe

is fixed in a predetermined recess.

Materials forming the matrix pattern include, for example, metals (chrome, aluminum, gold, etc.) and resins. They include resins such as acryl,  
5 polycarbonate, polystyrene, polyimide, acrylate monomers and urethane acrylate, and photosensitive resins such as photoresists having black dyes and black pigments contained therein. For specific examples of photosensitive resins, UV resists, DEEP-UV resists,  
10 ultraviolet cured resins and the like can be used. UV resists may include negative resists such as cyclized polyisoprene-aromatic diazide resists, phenol resin-aromatic azide compound resists, and positive resists such as novolac resin-diazonaphthoquinone resists.

15 DEEP-UV resists may include, for example, radiation dispersion type polymer resists such as polymethyl methacrylate, polymethylene sulfone, polyhexafluorobutyl methacrylate, polymethyl isoprobenil ketone and bromo poly 1-  
20 trimethylcylilpropine, and dissolution inhibiting resists such as cholate o-nitrobenzyl ester as positive type resists, and may include polyvinylphenol-3-3'-diazidediphenylsulfone, and polymethacrylate glycidyl as negative type resists.

25 Ultraviolet cured resins may include polyester acrylate, epoxy acrylate and urethane diacrylate containing approximately 2 to 10% by weight of one or



more types of photopolymerization initiators, which are selected from benzophenone and substituted derivatives thereof, oxime compounds such as benzyl, and so on.

For curbing reflection by the material forming the matrix during detection, light-blocking materials can be effectively used for materials forming the matrix pattern. For this purpose, it is effective to add black pigments in the above described resin, and for black pigments, carbon black and black organic pigments can be used.

Here, if the matrix 125 is composed of resin, the surface of the matrix 125 is hydrophobic. This structure is preferred when aqueous solution is used as a solution containing probes to be supplied to the well. That is, even if the probe solution is supplied to the well, the probe solution is supplied to a desired well quite smoothly. Also, if different probes are supplied among adjacent wells at a time, intermingling (cross contamination) of different probe solutions supplied among these wells can be prevented.

The thickness of the matrix (height from the solid surface) is determined in the light of matrix pattern forming process and the volume of the well, but it is preferably in the range of 1 to 20  $\mu\text{m}$ . Particularly, it can be considered as a thickness range allowing cross contamination to be prevented effectively when the probe solution is supplied to each well though an

ink jet process.

(Types of samples to be spotted)

Samples to be spotted as droplets onto the above described matrices of biological samples include probe  
5 nucleic acids, proteins and chemicals such as drugs.

For probe nucleic acids, in addition to deoxyribonucleic acid, any types of nucleic acids such as ribonucleic acid and peptide nucleic acid may be used as long as they have nucleic acid bases. The  
10 length of the oligonucleotide probe is not particularly limited, but it is preferably in the range of 10 mer to 50 mer for carrying out accurate hybridization reaction with cDNA.

For proteins, their own fluorescence can be used  
15 to detect DNA bonding proteins.

Some chemicals can also be detected with their own fluorescence.

(Method of preparing sample arrays)

Methods of spotting sample solution on defined  
20 positions in the size of several tens to several hundreds of microns include a pin system, an ink jet system and a capillary system.

The pin system refers to a method in which the sample is attached to the pin tip, for example, in such  
25 a manner that the pin tip is contacted with the surface of the solution including the sample, and then the tip is mechanically contacted with the solid phase, thereby

preparing a sample array. The capillary system using a capillary is such that the sample solution once sucked up to the capillary is mechanically contacted with the solid phase through the tip of the capillary as in the case of the pin system, thereby supplying the sample solution in an array form. For these spotting operations, various apparatuses commercially available from various companies may be used. These methods are considered as most preferable methods in the sense that any sample DNA can be supplied. However, as for quantification, the problem may be unsolved that viscosity varies depending on the length and concentration of DNA. For proteins, these methods are also preferred in the sense that they are deposited independently of the size and viscosity of molecules, but not suitable for quantitative analysis.

(Outline of sample array preparing methods through the ink jet process)

Samples capable of being discharged in an ink jet process include chemicals in addition to nucleic acids and proteins.

In the ink jet process, because shearing force is exerted, the length of dischargeable nucleic acids and the size of dischargeable proteins are often limited. However, it is superior in quantification to the pin system and capillary system, and is used more suitably than other systems with respect to discharge of

chemicals. Dischargeable nucleic acids are limited to those with relative length to bases of 5 kb or smaller, and dischargeable proteins are limited to those of 1000 K daltons or less. As for chemicals, any chemicals can be discharged.

For liquids for discharge to be used for discharging and supplying samples with ink jets, any liquid can be used as long as it is capable of being discharged from ink jets, and the above described liquid discharged from the head is shot in a predetermined position, and in the state of being mixed with nucleic acid probes and during discharge, the above described nucleic acid probes are not damaged.

And, in terms of dischargeability from the ink jet, particularly from the bubble jet head, for the properties of the above described liquid, it is preferable that its viscosity is in the range of 1 to 15 cps and its surface tension is 30 dyn/cm or larger. Also, if the viscosity is in the range of 1 to 5 cps and the surface tension is in the range of 30 to 50 dyn/cm, the position in which the liquid is spotted on the solid phase is extremely accurate, allowing the method to be used particularly suitably.

Therefore, if the stability of nucleic acid during discharge or the like is taken into consideration, a nucleic acid probe of, for example, 2 to 5000 mer, particularly 2 to 1000 mer is preferably contained in

the solution in concentrations of 0.05 to 500  $\mu\text{M}$ , particularly 2 to 50  $\mu\text{M}$ .

FIG. 3 is a schematic explanatory view of a specimen solution discharging method through the bubble jet process that is one embodiment of the present invention. In FIG. 3, reference numeral 101 denotes a liquid supplying system (nozzle) retaining a solution including a specimen as discharge liquid in such a manner that the solution is capable of being discharged, reference numeral 103 denotes a solid phase having a nucleic probe bound thereto with which the above described specimen is reacted, and reference numeral 105 denotes a bubble jet head having a function of giving heat energy to the above described liquid to discharge it, which is a type of ink jet head. Reference numeral 104 denotes a liquid including the specimen discharged from the bubble jet head. FIG. 4 is a 4-4 line sectional view of the bubble jet head 105 in FIG. 3, and in FIG. 4, reference numeral 105 denotes the bubble jet head, and reference numeral 107 denotes a liquid including a specimen solution to be discharged, and reference numeral 117 denotes a substrate portion having a heat generation portion to give discharge energy to the above described liquid. The substrate portion 117 includes a protective layer 109 formed by silicon oxide and the like, electrodes 111-1 and 111-2 formed by aluminum and the like, an

exothermic resistor layer 113 formed by nichrome and the like, a heat storage layer 115, and a support 116 formed by aluminum having good heat-release property.


5 The liquid 107 including the specimen comes to a discharge orifice (discharge outlet) 119, and forms a meniscus 121 with a predetermined pressure. Here, when electric signals are applied to the electrodes 111-1 and 111-2, a region (foaming region) denoted by reference numeral 123 abruptly releases heat, and the  
10 liquid 117 contacted therewith is discharged and flies toward the solid surface 103. The amount of liquid that can be discharged using a bubble jet head having such a structure varies depending on the size of its nozzle, but can be controlled approximately to 4 to 50  
15 picoliters, which is extremely useful as means for placing specimen probes in high density.

And, in terms of dischargeability from the ink jet, particularly from the bubble jet head, for the properties of the above described liquid, it is  
20 preferable that its viscosity is in the range of 1 to 15 cps and its surface tension is 30 dyn/cm or larger. Also, if the viscosity is in the range of 1 to 5 cps and the surface tension is in the range of 30 to 50 dyn/cm, the liquid is spotted in an exceedingly  
25 accurate position on the solid phase, allowing the method to be used particularly suitably.

Therefore, if the stability of nucleic acid during

discharge or the like is taken into consideration, a nucleic acid of, for example, 2 to 5000 mer, particularly 2 to 1000 mer is preferably contained in the solution in concentrations of 0.05 to 500  $\mu$ M, particularly 2 to 50  $\mu$ M.

For the composition of discharged liquid, the composition of liquid is not particularly limited, as long as the liquid has no substantial influence on the nucleic acid probe when it is mixed with the nucleic acid probe and when it is discharged from the ink jet, and it can be normally discharged to the solid phase using the ink jet, but preferable are liquids including glycerin, urea, thiodiglycol or ethylene glycol, isopropyl alcohol, and acetyl alcoholene expressed by the following formula.



(In the above formula (I), R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> represent alkyl groups, specifically linear or branched alkyl groups having 1 to 4 carbon atoms, m and n represent integer numbers, respectively, wherein m and n equal 0, or  $1 \leq m + n \leq 30$  holds, and if  $m + n = 1$  holds,

m or n equals 0).

Further specifically, a liquid containing 5 to 10% by weight (wt%) of urea, 5 to 10 wt% of glycerin, 5 to 10 wt% of thiodiglycol, and 0.02 to 5 wt%, more preferably 0.5 to 1 wt% of acetylene alcohol presented by the above formula (I) is suitably used.

The detecting method of the present invention is a method of detecting a complex formed between oligonucleotide for detection probes and an object component, which is used for the purpose of making evaluation/examination as to whether or not a component having capability of binding to oligonucleotide for use as a detection probe whose base sequence is known, and forming therewith a complex exists in a liquid test sample, and as to the degree of binding capability thereof if such a component exists in the sample. For detecting this complex, oligonucleotide for detection probes is fixed in advance on the solid surface substrate, whereby this fixed oligonucleotide is bound to the object component contained in the test sample, and the formed complex is separated while it is fixed on the solid substrate, and on the basis of a methodology for detecting complexes using proper detecting means, the amount of test samples required at this time is reduced to a very low level, and also, the detection accuracy and sensitiveness are kept at a sufficiently high level.



That is, in this methodology providing a base for the present invention, since surface density of the oligonucleotide for detection probes that is fixed on the solid surface substrate can be kept at a  
5 predetermined value, the amount of the formed complex is proportional to the binding capability of the object component, and is also proportional to the concentration of the object component contained in the test sample that is contacted with the solid surface  
10 substrate and is made to act on the oligonucleotide. Taking advantage of this characteristic, the test sample is contacted only with the surface with the oligonucleotide for detection probes actually fixed thereon, and the contact area is limited to a certain  
15 level, whereby the amount of used test sample is also limited to a certain level. Specifically, by adopting means for spotting a predetermined minimal amount of liquid in the form of droplets, the contact area and the amount of liquid put thereon is controlled with  
20 good reproducibility. The amount of the complex that would be automatically fixed on the solid surface substrate with formation is detected for this limited contact area, thereby achieving detection accuracy and sensitiveness essentially as high as those in the case  
25 of dipping in the liquid test sample the whole of the solid substrate with oligonucleotide for detection probes fixed thereon.

The complex is detected by the label bound on the surface of the substrate. When a complex of the oligonucleotide and the labeled test sample is formed and the individual spots are sufficiently spaced from each other, detection can be carried out independently for each spot. Therefore, if given or larger spaces are provided between adjacent spots, even though there are spots for different test samples nearby, only spots for desired test samples can be selected to continue detection work without being influenced by those spots. In the detection method of the present invention, in order to satisfy reliably this requirement that given or larger spaces be provided between adjacent spots, a defined array is formed in the spot position as a result of providing predetermined spaces as spaces between spots, and a predetermined amount of sample solution is spotted for each spot to make the spot area (contact area) constant, or make the spot diameter constant to ensure reproducibility because the shape of the spot (contact surface) is generally a circle. As a matter of course, for precluding influence of adjacent spots, a space between spots is selected such that optical signals (fluorescent) and the like derived from the adjacent spots are not mixed in the detection system, in the light of the measured area (diameter in the measurement range) of the detection system selected as appropriate in accordance with the spot diameter.

Also, as a matter of course, the detecting method of the present invention really shows its advantages in the case where there exist two or more types of test samples, and they are detected simultaneously.

5           On the other hand, on the surface to which a plurality of spots of such array forms is provided, one type of oligonucleotide for detection probes should be fixed in uniform surface density. Also, for the section in which the oligonucleotide for detection probes is fixed, its area and shape are selected as appropriate in accordance with the above described array space and the total number of spots to be included in a series of arrays. It is also possible to provide sections having different oligonucleotides fixed therein in different regions on the detection substrate to be used, and place a plurality of sections with two or more oligonucleotides fixed therein, respectively. That is, it can be said that the detecting method of the present invention becomes a more suitable method if used when two or more types of nucleotides are used as detection probes to carry out a series of evaluations simultaneously for a plurality of test samples, with respect to two or more types of object components corresponding to respective oligonucleotides.

10

15

20

25

Generally, in such an evaluation, it is often the case not that the oligonucleotides for detection probes

are predetermined while only an approximate number of test samples to be evaluated is determined. In such a case, it is preferable that as a detection substrate with oligonucleotides for detection probes fixed thereon in advance, a detection substrate with two or more types of detection probes put thereon systematically, having on the substrate surface in a matrix form sections in which respective oligonucleotides are fixed. In this detection substrate with fixed sections arranged thereon in a matrix form, the unit of total number of spots that are made in an array form in each section is fixed, but a plurality of these units of number of spots can be used to carry out evaluation depending on the number of test samples to be actually evaluated, thus enhancing convenience in practice. Furthermore, for the each section arranged in a matrix form, a pattern formed by hydrophobic compounds is preferably provided in its substrate to provide a form in which mutual regions are separated from one another.

In the detecting method of the present invention, nucleic acid molecules may be selected as object components to apply the same to evaluation as to whether or not they are engaged in double-strand formation into hybrid substances through hybridization reaction with oligonucleotide for detection probes. In this case, the method is an effective method in which

evaluation is made at a time even for multiple test samples, as to whether or not nucleic acid molecules including base sequences complementary to known base sequences that oligonucleotide for detection probes has  
5 are contained in the test sample. Alternatively, if two or more types of nucleotides for detection probes are provided, and one type of nucleic acid molecules are contained in each test sample, evaluation can be made for the nucleic acid molecule of which base  
10 sequences are still unknown, as to whether or not the nucleic acid molecule includes base sequences complementary to known base sequences that each oligonucleotide has, which is effective, for example, for means for making search for a gene group having a  
15 set of homologies.

The detection substrate of the present invention is a DNA probe substrate with oligonucleotides for use in probes respectively bound to sections arranged in a matrix form in advance, and particularly for the  
20 substrate itself, the bottoms of sections separated by wells (walls) of frame structure matrix patterns formed in advance by hydrophobic compounds are formed as hydrophilic surfaces, thereby making the binding of oligonucleotide easier. Also, by providing this  
25 hydrophobic wall, intermingling of DNA probes among adjacent sections can be curbed more reliably.

Also, using these DNA probe substrates, the test

sample is spotted in an array form on the matrix of oligonucleotide to carry out hybridization reaction, thereby providing means for checking quickly whether or not nucleic acid molecules having complementarity are included in each test sample for a certain oligonucleotide probe.

In this method, since the number of test samples that are used in hybridization reaction is determined depending solely on the number of spots, the size of the detection substrate is not limited, and by using a substrate of large area, the section in which each probe is fixed can be widened and necessity to enhance density can be eliminated. Thus, since the section in which each probe is fixed can be widened, a wide range of methods can be used such as methods in which a liquid containing probes is applied to, or printed as a "solid printed pattern" through ink jet process on defined regions on the substrate, or methods in which chemical synthesis is carried on the substrate, for means for binding of the probe/oligonucleotide on the substrate.

Also, considering that the probe/oligonucleotide is less expensive and easier to obtain than the test sample, no significant problem arises even if the area of the region to which oligonucleotide is bound is more or less increased, and in this case, with respect to a various kinds of test samples to be spotted, it is not

necessary to always spot them in high density.

Furthermore, when the test sample is spotted in small amounts, the concentration of the object component that is contained in the test sample is increased, whereby hybridization reaction can be accelerated, thus making it possible to perform high sensitive detection for a short time. In addition, application of the detecting method of the present invention will open the door to fields that could not be considered previously because a sufficient amount of samples could not be obtained, for example a new field in which mRNA obtained from tissues is directly examined.

Furthermore, information of reactivity in association with obtained hybridization reaction is analyzed/evaluated in terms of existence/not existence of complementarity to various kinds of oligonucleotides/probes, with respect to nucleic acid molecules contained in a specific test sample, thereby making it possible to carry out detection having also functions similar to those of conventional DNA arrays (hybridization reaction with multiple probes for one specimen).

Furthermore, the detecting method of the present invention provides means for evaluating as object components the interaction between chemicals, in particular drugs and oligonucleotides, the bonding of proteins to oligonucleotides and the like, and

therefore, it can also be used as means making it possible to examine object components included in the test sample for multiple items, with respect to a large number of test samples. In addition, it provides means  
5 making it possible to carry out examination on the same substrate at a time and under same conditions even for object components of different properties chemicals, proteins and nucleic acids.

The detecting method of the present invention and  
10 the detection substrate for use exclusively therein will be described further in detail below.

In FIG. 6 is shown an example of applying the detecting method of the present invention to an embodiment in which using cDNA as an object component,  
15 a hybrid substance is formed through hybridization reaction with oligonucleotide of known base sequence that is used for detection probes. In the detection substrate shown in FIG. 6, a plurality of rectangular sections separated systematically in a matrix form in advance is provided on the solid surface substrate of  
20 rectangular form. The rectangular sections are each spatially isolated by matrix compartments that are surrounding walls. DNA probes that are used for hybridization/probes are each bound uniformly to the  
25 bottom surface of the rectangular sections.

Also, attached is an enlarged view showing schematically a situation in which a plurality of test



samples including cDNA as an object component, for example two or more types of cDNA solutions prepared based on m-RNAs collected respectively are spotted in the form of the two-dimensional array like a square matrix, in a section with the DNA probe fixed therein. The detection substrate, the detection probe, components to be detected and the like that are used in the detecting method of the present invention will be described further in detail.

10           (Oligonucleotides that are used for detection probes)

          In the detecting method of the present invention, deoxyribonucleic acid can be used for oligonucleotide that is used for detection probes. In addition thereto, ribonucleic acid, peptide nucleic acid and the like can be used. Types thereof are not limited as long as they have desired base sequences, and are capable of being bound to other molecules in those portions, and also as long as they can be fixed on a solid substrate. Also, for portions excluding nucleic acid chains, those modified with non-nucleic acid atom groups and those having additional structures and so on can also be used as long as the above described requirements are satisfied.

25           Furthermore, for this oligonucleotide that is used for detection probes, a desired amount thereof should be artificially prepared or collected, and its base

sequence itself should be known. However, its nucleic acid part should have at least two bases. Its base length is not limited in principle, but if the length exceeds that of 100 bases, difficulty becomes more significant as its base length increases when fixation on the solid substrate is to be carried out, and therefore the base length is preferably restricted to that of 100 bases or less.

For example, when this oligonucleotide is subjected to hybridization reaction with, for example, nucleic acid molecules with the length of more than 100 bases, the length of the oligonucleotide is preferably at least 10 mer for obtaining sufficient bonding. On the other hand, if the length exceeds 50 mer, it is difficult to set conditions for controlling detection of mismatching, thus making it difficult to select and detect only those that are fully matched. Thus, in order to detect mutations, the length is preferably 60 mer or smaller.

Furthermore, the range of 10 mer to 60 mer is a preferable range even when oligonucleotide having desired base sequences, for example DNA is prepared through chemical synthesis.

(Shapes of sections with oligonucleotide fixed therein, which is arranged in a matrix form)

The shape of a section itself in which oligonucleotide for detection probes is bound and fixed

is not particularly limited. However, if considering that a test samples is spotted in an array form on this section, generally a simpler shape rather than a complicated outside shape is preferably selected. In addition, also when oligonucleotide is bound and fixed, generally a simpler shape is preferably selected for providing uniform surface density in such a section , in terms of working efficiency and convenience. Specifically, rectangular forms, for example, line forms, squares and rectangles are preferably adopted. Of course, in principle, forms whose perimeters are formed by curves such as circles and ellipses do not cause any problems.

On the other hand, in the detection substrate of the present invention, when two or more oligonucleotides that are used for detection probes are put on one substrate, sections in which they are fixed are preferably arranged in a matrix form, in terms of working efficiency and convenience. Also, preferably, the form of each section is unified, and its area is also unified.

(Density of sections arranged in a matrix form)

The density of sections arranged in a matrix form is selected as appropriate depending on the number of oligonucleotides that are put on the detection substrate at a time, but the density of 400 per centimeter square or less is preferable. If the

density is  $400/\text{cm}^2$ , and the form of each section is a square, the size of each section is a  $500\text{ }\mu\text{m}$  square. If test samples are closely arranged in an array form as spots with diameters of  $100\text{ }\mu\text{m}$ , 25 spots are arranged in total with 5 spots high by 5 spots wide. Also, if the diameter of the spot is  $20\text{ }\mu\text{m}$ , the number of spots that can be arranged in a row is 25, leading to 625 spots in total. Since the detecting method of the present invention has more significant advantages when there are a large number of test samples and they are examined at a time, the final object of the invention will be more satisfactorily achieved if the density of the section that is arranged is selected so that at least the upper limit of the number of test samples that can be spotted approximately equals the above described value.

For example, when the detecting method of the present invention is applied to test samples including cDNA, the number of test samples to be examined, specifically the total number of types of cDNA often is approximately as many as 3600. In this case, if the diameter of the spot is  $100\text{ }\mu\text{m}$ , the size of one section approximately equals a  $6\text{ mm}$  square when 60 spots are arranged in rows and columns, respectively. Also, even if the diameter of the spot is  $20\text{ }\mu\text{m}$ , the size of one section should be a  $1.2\text{ mm}$  square. In this way, in the detection substrate for use in the detecting method of

the present invention, there are not a few cases of application objects where the density of sections that are arranged in a matrix form is preferably selected as 400 per centimeter square or less.

5           Furthermore, in the detecting method of the present invention, the test sample is spotted as droplets, and in the case where the diameter of spot is 100  $\mu\text{m}$ , for example, the amount of liquid required for the droplet of one spot is about 25 picoliters. Even  
10 if the number of probes for use in examination is selected as 400 (for example, the number of sections of the matrix to be provided on the substrate is 400) for this spot size, the total amount of liquid required for the whole spots may be no more than 10 nanoliters for  
15 each test sample, thus making it possible to carry out objective examination items with a minimal amount of liquid.

          Also, in the conventional method in which the detection substrate is dipped in the solution of the  
20 test samples, the amount of required liquid is dependent on the size of the substrate, and thus if the amount of the test sample is essentially very small, the size of the substrate should be reduced in accordance with the amount of liquid, and it is  
25 essential to highly integrate probes that are fixed on the substrate. On the other hand, in the detecting method of the present invention, the size of the

substrate itself can be freely selected without allowing for the liquid amount the test sample. In addition, when oligonucleotide that is used for detection probes is fixed, the surface density should be uniformed as a matter of course, but it is not necessary to highly integrate a plurality of probes to fix them, thus making the fixing operation easier.

(Fixation of oligonucleotide on the substrate)

As means for fixing oligonucleotide that is used for detection probes on the surface of the substrate, a method in which oligonucleotide separately prepared in advance is supplied in predetermined sections by coating or printing to bind the oligonucleotide, or a method in which each oligonucleotide, specifically a DNA probe or the like is synthesized in solid phase on the substrate to prepare originally bound DNA can be used. Furthermore, even in the case where the oligonucleotide is not DNA but, for example, ribonucleic acid or peptide nucleic acid, synthesis on the substrate can be carried out to bind the oligonucleotide as described later.

On the other hand, when oligonucleotide, specifically DNA or ribonucleic acid, peptide nucleic acid or the like separately synthesized or collected in advance is used for detection probes, a process of fixing the oligonucleotide by covalent bond or of fixing it electrostatic coupling on the surface of the

substrate can be used.

(Synthesis of oligonucleotide on the substrate)

Synthesis of DNA on the substrate includes  
synthesis on the silicon substrate using  
5 photolithography as a methodology disclosed in U.S.  
Patent No. 5445934. The U.S. Patent No. 5445934 shows  
a method in which high density DNA probe arrays are  
prepared by dividing the surface of the silicon  
substrate into very small areas, and synthesizing DNA  
10 for probes. On the other hand, in the detection  
substrate for use in the detecting method of the  
present invention, for example, the size of the section  
in which each probe is fixed may be a 0.5 mm square or  
larger, and thus it is not always necessary to enhance  
15 density. However, also in the detection substrate of  
the present invention, photodecomposable protective  
groups, protective groups that are decomposed by  
chemicals and the like are bound to nucleic acid in  
advance, and processes of masking, light exposure and  
20 reaction are repeated, whereby DNA chains can be  
synthesized on each section using the methodology  
described in the U.S. Patent Publication No. 5445934 in  
which four types of nucleic acid bases are bound for  
each base to stretch the DNA chain having desired base  
25 sequences.

(Fixation of oligonucleotide synthesized or  
collected in advance)

As means for carrying out fixation using electrostatic coupling, a method in which polylysine, polyethyleneimine and polyalkylamineaone on the solid surface substrate are subjected to blocking using the  
5 negative charge of DNA is generally used.

However, in the case of oligonucleotide with base length of 60 or less that is not sufficiently long, the electric charge of its phosphate groups is also weak, and thus binding onto the substrate by the above  
10 described method is not necessarily strong. For this oligonucleotide whose base length is not sufficiently long, if a method in which oligonucleotide with functional groups for covalent bond introduced in the terminal of nucleic acid is synthesized in advance, the  
15 substrate is subjected to surface processing suitable for functional groups, and the above described functional groups are used to accomplish covalent bond is used, stronger binding can be achieved, which is more preferable.

20 Also, in the case where the oligonucleotide is RNA, the above described method that is used for DNA may be applied. Alternatively, in the case where the oligonucleotide is peptide nucleic acid, its nucleic acid part may be used to apply the above described  
25 method that is used for DNA.


(Types of functional groups for use in fixation by covalent bond between the solid substrate and



oligonucleotide)

When oligonucleotide is fixed on the solid surface substrate through covalent bond, functional groups are generally introduced in oligonucleotide and the solid surface substrate, respectively, in advance to carry out reaction therebetween. For this combination of functions, a preferable example is a combination such that maleimide groups are introduced in the surface of the substrate and thiol groups (-SH) are introduced in oligonucleotide. Specifically, thiol groups (-SH) are bound to the terminal of oligonucleotide while the solid surface is subjected to processing of forming a coating having maleimide groups, and when oligonucleotide is supplied to the solid surface, the thiol groups (-SH) are made to act on and react with the maleimide groups to perform fixation through formation of covalent bond.

For introducing maleimide groups in the solid surface, various kinds of methods may be used, and for example, an aminosilane coupling agent is reacted with a glass substrate, and then a reagent (EMCS reagent: manufactured by Dojin Co., Ltd.) including N-(6-maleimidocaproyloxy) succinimide expressed by the following formula, whereby a coating layer having maleimide groups can be formed.



5

For another example, a reagent containing succinimidyl 4-(maleimidophenyl)butyrate can be used to react with amino groups preferably.

Also, for example, oligonucleotide with thiol groups introduced therein can be synthesized by using 5' -Thiol-Modifier C6 (manufactured by Glen Research Co., Ltd.) as a five prime-end reagent when DNA is synthesized using a DNA automatic synthesizing apparatus. Furthermore, after synthesis, purification processing by high speed liquid chromatography is applied after normal deprotection reaction.

Combinations of functional groups capable of being used for fixation by covalent bond include, for example, a combination of epoxy groups (on the solid surface) and amino groups (the terminal of oligonucleotide) in addition to the above described combination of thiol groups and maleimide groups. Methods for introducing epoxy groups in the solid surface include, for example, a method in which coating is applied to the solid surface constituted by polyglycidyl methacrylate having epoxy groups and a method in which a silane coupling agent having epoxy

groups is applied to the solid surface made of glass and is reacted with glass.

(Supply of oligonucleotide solution by the ink jet process)

5           There is no particular limitation on means for supplying predetermined sections on the surface of the solid substrate with a solution containing oligonucleotide to be fixed thereon, as long as a uniform amount of liquid is supplied for each unit  
10   area. In the case where printing by the ink jet process and the like is used, a "solid print pattern" is prepared, and then using an ink jet type printer head that is used for ink jet printers, the cartridge for the ink is filled with oligonucleotide solution  
15   instead of the ink, and printing for a defined area is carried out. If the amount of liquid to be supplied is small, items of large volume like an ink cartridge are not used, and instead a structure in which a sample supplying portion such as a tube is connected to a head  
20   to supply the oligonucleotide solution to the head may be used.

          For the oligonucleotide solution for discharge, which is used in this method, a solution that is capable of being discharged in the form of ink jets,  
25   and has viscosity suitable for a minimal amount of droplets discharged from the head to be shot onto a desired position is used. In addition, a solvent to be


used is selected from solvents that satisfy the above described requirements and give no damages to desired oligonucleotide in the state of being mixed with the desired oligonucleotide and during discharge.

5           Specifically, in terms of dischargeability from the ink jet head, particularly from the bubble jet head, it is preferable that for example the viscosity is in the range of 1 to 15 cps, and the surface tension is 30  
10           dyn/cm or larger as the properties of the solution. In particular, when the viscosity being in the range of 1 to 5 cps and the surface tension being in the range of 30 to 50 dyn/cm are selected, the position in which the solution is shot onto the substrate is extremely accurate, and a supplying method using the bubble jet  
15           head is particularly suitably used.

          In addition, if stability of oligonucleotide during discharge and the like are considered, the supplying means of the ink jet system is further preferred when for example, a solution containing  
20           oligonucleotide of 2 to 100 mer, particularly of 2 to 60 mer in concentrations ranging from 0.05 to 500  $\mu$ M, preferably from 2 to 50  $\mu$ M is used.

          In applying a discharging method of the ink jet system, the liquid composition of the oligonucleotide  
25           solution is not particularly limited, as long as the solution gives practically no damages to desired oligonucleotide in the state of being mixed with the

desired oligonucleotide and during discharge as a matter of course as described above, and it can be discharged to the surface of the solid substrate using the ink jet. Furthermore, preferable is a solution containing, for example, glycerin, urea, thiodiglycol or ethylene glycol, isopropyl alcohol, or acetylene alcohol expressed by the following formula in addition to desired oligonucleotide.



(In the above formula, R1, R2, R3 and R4 represent alkyl groups, for example linear or branched alkyl groups having 1 to 4 carbon atoms, respectively, and m and n represent 0 or positive integer numbers, respectively, and satisfy  $1 \leq m + n \leq 30$ ). In addition, specifically the liquid composition including 5 to 10 wt% of urea, 5 to 10 wt% of glycerin, 5 to 10 wt% of thioglycol, and 0.02 to 5 wt%, more preferably 0.5 to 1 wt% of acetylene alcohol expressed by the formula (I) allows the discharging method of the ink jet system to be used suitably.

(Structure of matrices composed of hydrophobic

walls and hydrophilic wells)

Also, for sections of matrix form that are provided on the solid surface, for example, sections of matrix form constituted by hydrophobic walls (barriers) surrounding hydrophilic wells (recesses) may be formed to prevent coupling between adjacent sections. A structure may also be used in which the solution of oligonucleotide is supplied to the hydrophilic wells (recesses) surrounded by the hydrophobic walls (barriers), and oligonucleotide is fixed only in the bottom of the hydrophilic wells (recesses).

(Materials of walls/wells)

When as sections arranged in a matrix form, the solution of oligonucleotide is supplied to the bottom of the wells (recesses) separated by wall (barrier) patterns to carry out binding reaction, it is desirable that the bottom of the wells (recesses) is wetted densely with the solution, but the walls (barriers) have poor wettability with the solution. For example, it is preferable that the solid material constituting the surface of the bottom of the wells (recesses) is much hydrophilic, and the surface of the walls (barriers) and the portion corresponding to partitions with neighboring sections are less hydrophilic. The solution of oligonucleotide supplied in the bottom of the well (recess) is spread across the bottom, but is prevented from finding its way over the wall (barrier)

into adjacent sections. Also, even the droplet erroneously supplied in the position related to the wall (barrier) quickly moves into a desired well (recess) having good wettability, and as a result a predetermined amount of oligonucleotide solution can be supplied in the well (recess) more reliably.

An example of sections arranged in a matrix form that is provided on the detection substrate of the present invention is shown in FIG. 10. The sections in a square matrix form have a structure in which heights (walls) having frame structures are provided on the surface of the solid substrate, and arranged rectangular recesses (wells) are separated. Specifically, the recesses (wells) separated from one another by the heights (walls) having frame structures are formed by coating the entire surface of the solid substrate with a material forming heights (walls), and thereafter providing rectangular through-holes (cut-off portions) to open recesses (wells). Thus, the bottom of the recess (well) has an exposed surface of the solid substrate. The exposed portion of the surface of the solid substrate is subjected to processing for providing a surface to which oligonucleotide can be bound. As a result, oligonucleotide is fixed only in the bottom of this recess (well).

Materials forming heights (walls) having frame structures include, for example, metals (chrome,

aluminum, gold, etc.) and resins. Resins include resins such as acryl, polycarbonate, polystyrene, polyimide, acrylate monomers and urethane acrylate, and photosensitive resins such as photoresists having black dies and black pigments contained therein.

Furthermore, for specific examples of photosensitive resins, UV resists, DEEP-UV resists, ultraviolet cured resins and the like can be used. UV resists may include negative resists such as cyclized polyisoprene-aromatic pisazide resists, phenol resin-aromatic azide compound resists, and positive resists such as novolac resin-diazonaphthoquinone resists.

DEEP-UV resists may include, for example, radiation dispersion type polymer resists such as polymethyl methacrylate, polymethylene sulfone, polyhexafluorobutyl methacrylate, polymethyl isoprobenil ketone and bromo poly 1-trimethylcylilpropine, and dissolution inhibiting resists such as cholate o-nitrobenzyl ester as positive type resists, and may include borovinylphenol-3-3'-diazidediphenylsulfone, and polymethacrylate glycidyl as negative type resists.

Ultraviolet cured resins may include polyester acrylate, epoxy acrylate and urethane diacrylate containing approximately 2 to 10% by weight of one or more types of photopolymerization initiators, which are selected from benzophenone and substituted derivatives



thereof, oxime compounds such as benzyl, and so on.

When detection is carried out using a fluorescent mark, a light-blocking material can be used effectively for curbing light reflex by the material forming this height (wall) having a frame structure. For providing a light-blocking property, it is effective to add black pigments in the above described resins, and in this case, black pigments that can be used may include carbon black and black organic pigments.

Furthermore, if the height (wall) having a frame structure is formed by the above described hydrophobic resin, the surface of the height (wall) is hydrophobic. The configuration in which heights (walls) having frame structures that are formed by hydrophobic materials are provided is more preferable in the case where an aqueous solution is used as a solution containing oligonucleotide to be supplied to the surface of the substrate of recesses (wells). Even if the aqueous solution is supplied in a position related to the surface of the height (wall), it is not persistently attached to the surface of the wall, but gradually moves to the bottom of the recess (well) located in a lower position. Also, solutions of different oligonucleotides are supplied to adjacent recesses (wells), but they are separated from each other by the hydrophobic height (wall), and therefore intermingling (cross contamination) between solutions due to

penetration of liquid is prevented.

Furthermore, for the thickness (height from the solid surface) of the height (wall) having a frame structure, the volume of the recess (well) is selected in the light of the amount of the oligonucleotide solution that is supplied to the recess (well), and the thickness is determined as appropriate so that the volume is filled with the solution. Also, depending on methods of forming the height (wall), the thickness is preferable selected such that it is in the range of 1 to 20  $\mu\text{m}$  and satisfies the above described requirement. The thickness of the height (wall) selected in this way is in the range of thickness allowing cross contamination between adjacent wells to be prevented effectively when the oligonucleotide solution is supplied to each well by the ink jet process.

(Types of specimens)

Object components contained in the test sample to which the detecting method of the invention can be applied include mRNA, cDNA, proteins, cell extracts and chemicals such as drugs.

Furthermore, when cDNA is used as an object component, it is possible to use double-strand cDNA directly, but the single-strand cDNA marked in advance is preferable in forming hybrid substances efficiently and performing detection thereof conveniently.

On the other hand, mRNA is of essentially single-

strand, and it is marked in some way to form marked mRNA, thereby making it possible to form hybrid substances efficiently and perform detection thereof. Furthermore, the amount of mRNA in the test sample is  
5 generally small, and it is an object component more remarkably reflecting the advantage that the amount of sample solution required for detection can be reduced to a low level, which is characteristic of the detecting method of the present invention. However,  
10 since admixture of RAN decomposition enzymes tends to occur during handling, a predetermined amount of substance to curb decomposition of mRNA such as RNA decomposition enzyme inhibitors such as diethyl pyrocarbonate is desirable added in the test sample  
15 solution. In addition to mRNA, similarly, the genome of RNA viruses can be an object component. In addition, tRNA, ribosomal RNA and the like can be object components.

On the other hand, when the protein is used as an  
20 object component, formed complexes can be detected using the fluorescence emitted by the protein itself.

Also, some chemicals also emit their own fluorescence, and enables formed complexes to be detected using the fluorescence. Chemicals that do not  
25 emit fluorescence may be marked by methods using functional groups of compounds. Those to which the detecting method of the invention can be applied may

include, for example, chemicals that can be bound to single-strain DNA. In addition, they may include, for example, chemicals that can be bound to single-strain RNA.

5 (Means for spotting test samples in an array form)

In the detecting method of the present invention, the test sample is spotted in an array form in a defined position on the detection substrate. For the purpose of reducing amount of required liquid to a  
10 minimal level, the spot diameter is selected so that it is in the range of several tens to 100  $\mu\text{m}$ , but with such a spot diameter, the liquid should be spotted in high uniformity of spotted amounts and high positional accuracy. As means for satisfying this requirement,  
15 there are spotting apparatuses of pin systems, ink jet systems and capillary systems.

The pin system refers to a method in which the test sample is attached to the pin tip, and the end point thereof is mechanically contacted with the solid  
20 surface, thereby taking out a fixed amount of the test sample. The capillary system using capillaries refers to a method in which the test sample solution is sucked up to the capillary on a temporary basis, and the tip of the capillary is mechanically contacted with the  
25 solid surface as in the case of the pin system, thereby taking out a fixed amount of the test sample. A various kinds of spotting apparatuses adopting these

two systems are commercially available, and thus commercially available apparatuses may be used.

5 The spotting apparatuses of the pin system and capillary system enable any types of test samples to be spotted, and are considered as most preferable methods for unknown test samples. For example, however, the viscosity of the test sample solution is varied depending on the length and the concentration of DNA contained in the test sample, and therefore the amount of spotted liquid is varied. Thus, a problem arises in terms of quantification. Also with respect to proteins, the viscosity of the test sample solution is varied depending on the size of the molecules and the concentration, thus raising a problem in terms of quantification.

10

15

(Spots in an array form of test samples by the ink jet process)

Specimens that can be discharged by the ink jet process include chemicals in addition to nucleic acids and proteins.

20

In the ink jet process, because shearing force is exerted, the length of nucleic acids and the size of proteins that can be discharged are limited. However, it is superior in quantification to the pin system and capillary system, and is used more suitably than other systems, particularly with respect to discharge of chemicals. Preferably, dischargeable nucleic acids are

25

those with relative length to bases of 5 kb or smaller,  
and dischargeable proteins are those of 1000 K daltons  
or less. As for chemicals, their molecular weights are  
generally small enough compared to nucleic acids and  
5 proteins, and therefore any chemicals can be discharged  
except for polymers having extremely large molecular  
weights.

FIG. 3 illustrates schematically a method of  
discharging specimen solution by the ink jet process,  
10 particularly the bubble jet process, which is one means  
that is used for spotting test sample solution in the  
present invention. In FIG. 3, reference numeral 101  
denotes a liquid supply system (nozzle) retaining a  
solution including a specimen as discharge liquid in  
15 such a manner that the solution is capable of being  
discharged, reference numeral 103 denotes a solid phase  
having a nucleic probe bound thereto with which the  
specimen is reacted, and reference numeral 105 denotes  
a bubble jet head having a function of giving heat  
20 energy to the liquid to discharge it, which is a type  
of ink jet head. Reference numeral 104 denotes a  
liquid (droplet) including the specimen discharged from  
the bubble jet head. FIG. 4 is a sectional view of the  
bubble jet head 105 described in FIG. 3. In FIG. 4,  
25 reference numeral 107 denotes a liquid including a  
specimen solution to be discharged from the bubble jet  
head 105, and reference numeral 118 denotes a substrate

portion having a heat generation portion to give discharge energy to the above described liquid. The substrate portion 118 includes a protective layer 109 formed by silicon oxide and the like, electrodes 111-1 and 111-2 formed by aluminum and the like, an exothermic resistor layer 113 formed by nichrome and the like, a heat storage layer 115, and a support 116 formed by aluminum having good heat-release property. The liquid 107 including the specimen comes to a discharge orifice (discharge outlet) 119, and forms a meniscus 121 with a predetermined pressure. In this situation, when electric signals are applied to the electrodes 111-1 and 111-2, a region (foaming region) denoted by reference numeral 123 abruptly releases heat, and the liquid 117 contacted therewith is discharged and flies toward the solid surface 103. The amount of liquid that can be discharged using a bubble jet head having such a structure varies depending on the size of its nozzle, but can be controlled approximately to 4 to 50 picoliters, which is extremely useful as means for arranging probes in high density in a matrix form on the surface of the substrate.

And, in terms of dischargeability from the ink jet, particularly from the bubble jet head, for the properties of the above described liquid, it is preferable that its viscosity is in the range of 1 to 15 cps and its surface tension is 30 dyn/cm or larger.


Also, if the viscosity is in the range of 1 to 5 cps and the surface tension is in the range of 30 to 50 dyn/cm, the position in which the droplet is spotted (spot position) on the solid phase is extremely accurate, allowing the method to be used particularly suitably.

In addition, if the stability of nucleic acid during discharge or the like is taken into consideration, single-strain nucleic acid or double-strain nucleic acid of, for example, 2 to 5000 mer, particularly 2 to 10000 mer is preferably contained in the solution. For example, c-DNA chips are preferably contained in the concentration of 0.05 to 500  $\mu$ M, particularly 2 to 50  $\mu$ M.

For the composition of discharged liquid, the composition of liquid is not particularly limited, as long as the liquid has no substantial influence on the nucleic acid probe when it is mixed with the nucleic acid probe and when it is discharged from the ink jet, and it can be normally discharged to the solid phase using the ink jet, but preferable are liquids including glycerin, urea, thiodiglycol or ethylene glycol, isopropyl alcohol, and acetyl alcohol expressed by the following formula.



5



(In the above formula, R1, R2, R3 and R4 represent alkyl groups, specifically linear or branched alkyl groups having 1 to 4 carbon atoms, m and n represent 0 or positive integer numbers, respectively, and satisfy  $1 \leq m + n \leq 30$  holds).

Further specifically, a liquid containing 5 to 10% by weight (wt%) of urea, 5 to 10 wt% of glycerin, 5 to 10 wt% of thiodiglycol, and 0.02 to 5 wt%, more preferably 0.5 to 1 wt% of acetylene alcohol is suitably used.

#### [Examples]

The present invention will be described in detail below using Examples. Furthermore, the Examples shown herein represent one example of most suitable embodiments of the present invention, but the invention should not be limited by these Examples.

#### Example 1

A glass substrate with black matrices for specimen matrices for analyzing sequences of p 53 genes on a specimen matrix substrate partitioned by patterns is

prepared.

1. Preparation of a black matrix introduction substrate coated with polylysine. A glass substrate (60 mm x 50 mm) made of synthetic quartz is subjected to supersonic cleaning using 2% sodium hydrate solution, and is then subjected to UV ozone processing to clean the surface. Then, a polylysine solution (manufactured by sigma Co., Ltd.) is applied to the entire surface with a spin coater. In addition, a DEEP-UV resist (negative type resist for black matrices) (BK-739P manufactured by Nippon Steel Chemical Co., Ltd.) is applied thereto with the spin coater so that the thickness after curing is 5  $\mu$ m, and this substrate is heated for curing at 80°C for 5 minutes with a hotplate. Using a DEEP-UV aligner, a region of 1 cm x 1 cm is proximately exposed to light using a patterned mask so that the distance (X) between adjacent wells in FIG. 1 is 100  $\mu$ m and the form of the well is a square of 1 mm x 1 mm, and then development is carried out with a developing solution of inorganic alkaline solution using a spin drier, and the developing solution is washed out completely with purified water.

Then, the substrate is briefly dried using the spin drier, and is thereafter heated at 180°C for 30 minutes in a clean oven to have resist fully cured to obtain a substrate in which 400 wells are arranged as a

predetermined arrangement and adjacent wells are separated from each other by the black matrix. Furthermore, the volume of each well is calculated as 5  $\mu$ l if the thickness of the liquid is 5  $\mu$ m.

5           2. Fixation of specimen DNA

          (1) Preparation of cDNA libraries

The p 53 gene is obtained by a PCR reaction from 64 types of cDNA libraries obtained from tumor tissues.

66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225  
2226  
2227  
2228  
2229  
2230  
2231  
2232  
2233  
2234  
2235  
2236  
2237  
2238  
2239  
2240  
2

The PCR cycle is such that after thermal denaturation at 95°C for 5 minutes, cycles at 95°C for 30 seconds, at 55°C for 30 seconds and at 72°C for 60 seconds are conducted 29 times, and finally the solution is left for reaction at 72°C for 5 minutes and is then stored at 4°C.

After the reaction, gel electrophoresis is performed to confirm a product existing in the region of molecular weight of about 300 mer, and purification is carried out with MicroSpin Column S200 (Pharmacia) to obtain p 53 genes (p 53 DNA).

### (3) Synthesis of single-strain p 53 DNA

Using as a matrix the DNA obtained in the above (2), a single-strain marked DNA is obtained by the PCR reaction using 5' primer (Takara Shuzo Co., Ltd.). The composition of the reaction solution comprises

One shot LA PCR Mix	25 µl
5' primer (20 µM)	1
P 53 DNA	1
DW	22/50 µl, and

the reaction cycle is such that cycles at 96°C for 30 seconds, at 60°C for 15 seconds and at 60°C for 4 minutes are repeated 24 times, and finally the solution is stored at 4°C. Thereafter, it is purified with MicroSpin Column S200.

### (4) Fixation of p 53 cDNA

5 µl of the single-strain DNA obtained in the

above (3) is injected under a microscope into each well of the polylysine-coated substrate with black matrices prepared in the above (1), and is fixed through electrostatic coupling.

5           3. Analysis of variation of p 53 genes with oligonucleotide probes

10           The 64 DNAs were selected, focusing attention on the 248th and 249th amino acid sequences of the p 53 gene being a tumor inhibitor gene. That is, it is known that a case of frequent variation in the base sequence of CGGAGG is the case where the first C is changed to T, the second A is changed to G, and the third G of the sequence corresponding to the 249th amino acid is changed to T. Thus, the 64 probes are  
15           designed, focusing attention on the base sequence at these three points.

20           That is, it is a structure in which the total length of the probe is 18 mer, and six bases including this variation are located at the center thereof, and the bases are sandwiched between common sequences. A common sequence corresponds to the range from the five prime-end to the ATGAAC, and the subsequent portion including variation corresponds to the NNGAGN and a further subsequent common portion corresponds to the  
25           CCCATC, resulting in a final sequence of 5'ATGAACNNGAGNCCCATC3'. Here, the portion expressed by N corresponds to the A, G, C, and T that are four types

Country	Year	Population	Area	Population Density	Area Density
Algeria	1960	10,000,000	2,381,472	420	168
Algeria	1970	12,000,000	2,381,472	504	211
Algeria	1980	14,000,000	2,381,472	588	248
Algeria	1990	16,000,000	2,381,472	672	287
Algeria	2000	18,000,000	2,381,472	756	326
Algeria	2010	20,000,000	2,381,472	840	365
Algeria	2020	22,000,000	2,381,472	924	404
Algeria	2030	24,000,000	2,381,472	1,008	443
Algeria	2040	26,000,000	2,381,472	1,092	482
Algeria	2050	28,000,000	2,381,472	1,176	521
Algeria	2060	30,000,000	2,381,472	1,260	560
Algeria	2070	32,000,000	2,381,472	1,344	599
Algeria	2080	34,000,000	2,381,472	1,428	638
Algeria	2090	36,000,000	2,381,472	1,512	677
Algeria	2100	38,000,000	2,381,472	1,596	716
Algeria	2110	40,000,000	2,381,472	1,680	755
Algeria	2120	42,000,000	2,381,472	1,764	794
Algeria	2130	44,000,000	2,381,472	1,848	833
Algeria	2140	46,000,000	2,381,472	1,932	872
Algeria	2150	48,000,000	2,381,472	2,016	911
Algeria	2160	50,000,000	2,381,472	2,100	950
Algeria	2170	52,000,000	2,381,472	2,184	989
Algeria	2180	54,000,000	2,381,472	2,268	1,028
Algeria	2190	56,000,000	2,381,472	2,352	1,067
Algeria	2200	58,000,000	2,381,472	2,436	1,106
Algeria	2210	60,000,000	2,381,472	2,520	1,145
Algeria	2220	62,000,000	2,381,472	2,604	1,184
Algeria	2230	64,000,000	2,381,472	2,688	1,223
Algeria	2240	66,000,000	2,381,472	2,772	1,262
Algeria	2250	68,000,000	2,381,472	2,856	1,301
Algeria	2260	70,000,000	2,381,472	2,940	1,340
Algeria	2270	72,000,000	2,381,472	3,024	1,379
Algeria	2280	74,000,000	2,381,472	3,108	1,418
Algeria	2290	76,000,000	2,381,472	3,192	1,457
Algeria	2300	78,000,000	2,381,472	3,276	1,496
Algeria	2310	80,000,000	2,381,472	3,360	1,535
Algeria	2320	82,000,000	2,381,472	3,444	1,574
Algeria	2330	84,000,000	2,381,472	3,528	1,613
Algeria	2340	86,000,000	2,381,472	3,612	1,652
Algeria	2350	88,000,000	2,381,472	3,696	1,691
Algeria	2360	90,000,000	2,381,472	3,780	1,730
Algeria	2370	92,000,000	2,381,472	3,864	1,769
Algeria	2380	94,000,000	2,381,472	3,948	1,808
Algeria	2390	96,000,000	2,381,472	4,032	1,847
Algeria	2400	98,000,000	2,381,472	4,116	1,886
Algeria	2410	100,000,000	2,381,472	4,200	1,925
Algeria	2420	102,000,000	2,381,472	4,284	1,964
Algeria	2430	104,000,000	2,381,472	4,368	2,003
Algeria	2440	106,000,000	2,381,472	4,452	2,042
Algeria	2450	108,000,000	2,381,472	4,536	2,081
Algeria	2460	110,000,000	2,381,472	4,620	2,120
Algeria	2470	112,000,000	2,381,472		

Table 1

SEQ ID NO.	Sequence	SEQ ID NO.	Sequence
1	GATGGGACTCAAGTTCAT	33	GATGGGCCTCAAGTTCAT
2	GATGGGACTCAGGTTCAT	34	GATGGGCCTCAGGTTCAT
3	GATGGGACTCACGTTCAT	35	GATGGGCCTCACGTTCAT
4	GATGGGACTCATGTTCAT	36	GATGGGCCTCATGTTCAT
5	GATGGGACTCGAGTTCAT	37	GATGGGCCTCGAGTTCAT
6	GATGGGACTCGGGTTCAT	38	GATGGGCCTCGGGTTCAT
7	GATGGGACTCGCGTTCAT	39	GATGGGCCTCGCGTTCAT
8	GATGGGACTCGTGTTCAT	40	GATGGGCCTCGTGTTCAT
9	GATGGGACTCCAGTTCAT	41	GATGGGCCTCCAGTTCAT
10	GATGGGACTCCGTTCAT	42	GATGGGCCTCCGTTCAT
11	GATGGGACTCCCGTTCAT	43	GATGGGCCTCCCGTTCAT
12	GATGGGACTCCTGTTCAT	44	GATGGGCCTCCTGTTCAT
13	GATGGGACTCTAGTTCAT	45	GATGGGCCTCTAGTTCAT
14	GATGGGACTCTGGTTCAT	46	GATGGGCCTCTGGTTCAT
15	GATGGGACTCTCGTTCAT	47	GATGGGCCTCTCGTTCAT
16	GATGGGACTCTTGTTCAT	48	GATGGGCCTCTTGTTCAT
17	GATGGGGCTCAAGTTCAT	49	GATGGGTCTCAAGTTCAT
18	GATGGGGCTCAGGTTCAT	50	GATGGGTCTCAGGTTCAT
19	GATGGGGCTCACGTTCAT	51	GATGGGTCTCACGTTCAT
20	GATGGGGCTCATGTTCAT	52	GATGGGTCTCATGTTCAT
21	GATGGGGCTCGAGTTCAT	53	GATGGGTCTCGAGTTCAT
22	GATGGGGCTCGGGTTCAT	54	GATGGGTCTCGGGTTCAT
23	GATGGGGCTCGCGTTCAT	55	GATGGGTCTCGCGTTCAT
24	GATGGGGCTCGTGTTCAT	56	GATGGGTCTCGTGTTCAT
25	GATGGGGCTCCAGTTCAT	57	GATGGGTCTCCAGTTCAT
26	GATGGGGCTCCGTTCAT	58	GATGGGTCTCCGTTCAT
27	GATGGGGCTCCCGTTCAT	59	GATGGGTCTCCCGTTCAT
28	GATGGGGCTCCTGTTCAT	60	GATGGGTCTCCTGTTCAT
29	GATGGGGCTCTAGTTCAT	61	GATGGGTCTCTAGTTCAT
30	GATGGGGCTCTGGTTCAT	62	GATGGGTCTCTGGTTCAT
31	GATGGGGCTCTCGTTCAT	63	GATGGGTCTCTCGTTCAT
32	GATGGGGCTCTTGTTCAT	64	GATGGGTCTCTTGTTCAT

Then, for each of the 64 types of marked probe DNAs, a 8  $\mu$ M solution containing glycerin, urea and thiodiglycol in the final concentration of 7.5%, and acetylenol EH in the final concentration of 1% is

5 prepared. A different probe solution is charged by 100  $\mu$ l in each of the six nozzles of BJ Printer Head BC 62 (manufactured by Cannon Inc.). Arrangement is made so that six DNAs can be discharged for each head, and two heads are used to discharge 12 DNAs at a time, and the  
10 heads are exchanged 6 times to discharge DNAs so that each spot of 64 DNA is formed independently. In this way, total 64 probes are discharged in the form of the 8  $\times$  8 array in each well of black matrix coated with polylysine.

15 FIG. 5 shows an arrangement on each black matrix of 64 DNA probes that are discharged. In this case, 64 DNA probes are spotted in one matrix.

Thereafter, this substrate in which each probe is spotted is left in a humidifier chamber set at 40°C to  
20 carry out a hybridization reaction.

Thereafter, the substrate is cleaned with a 10 mM phosphate buffer containing 100 mM NaCl to remove DNA probes that have not been engaged in the formation of the hybrid substance.

25 DNA arrays after the hybridization reaction are observed using an inverted fluorescence microscope equipped with a filter set suitable for rhodamine.



If the gene as a specimen has normal base sequences, spots of highest fluorescence intensity should be observed in the gene at the location of the relative 42th DNA prove. It can be considered that those are derived from the hybrid of the p 53 gene having normal sequences amplified with the probe DNA and PCR. In a varied gene, detectable spots are observed at the location other than the 42th location, and a varied sequence can be known from the DNA probe supplied to the location.

#### Example 2

(Evaluation of existence/not existence of carcinogenic genes using mRNA)

##### 1. Extraction of mRNA

"QuickPrep Micro mRNA Purification Kit (manufactured by Amersham Pharmacia biotech co., Ltd.) is used to extract mRNA from tumor tissues collected with the biopsy. This mRNA is bound to a polylysine substrate with black matrices as in the case of Example

##### 1.

2. Examination of existence/not existence of carcinogenic genes and the type thereof with various kinds of carcinogenic gene probe arrays.

Sets of cloned oncogenes (18 types, manufactured by Takara Shuzo Co., Ltd.) are purchased, and then "LabelITnon-RI Labeling Kits" are used to perform rhodamine marking.

18 types of marked oncogene probes are spotted as an arrangement of 4 x 5 on the above described substrate with mRNA bound thereto, using a microarray preparing apparatus (pin system) manufactured by Cartesian Technologies Co., Ltd.

Further, a hybridization reaction is carried out as in the case of Example 1.

The type of oncogenes existing in the mRNA section extracted from each tissue can be known.

At this time, detection can be sufficiently performed with one type of marks irrespective of types of oncogenes.

The second invention will be describe more specifically with Examples below.

(Example 3)

An example of procedures for preparing an substrate with oligonucleotide bound thereto will be described below. In this embodiment, a detection substrate with oligonucleotide bound to a region of 2 mm square on a glass substrate was prepared in accordance with the procedure described below.

1. Cleaning of the substrate

A glass substrate of 1-inch square was placed in a rack, and was soaked in a detergent for ultrasonic cleaning. Thereafter, it was subjected to ultrasonic cleaning in the above described detergent for 20 minutes, followed by removing the detergent by rinsing.

Furthermore, it was rinsed with distilled water, followed by further performing ultrasonication for 20 minutes in a container containing distilled water.

Then, this glass substrate was soaked for 10 minutes in 1N sodium hydrate solution heated in advance. After it was taken out from the solution, the 1N sodium hydrate solution adhered to the surface was washed out with water, and thereafter cleaning with distilled water was continued.

## 10            2. Surface treatment

The above described cleaned glass substrate was soaked in an aqueous solution of 1% silane coupling agent (manufactured by Shin-Etsu Chemical Co., Ltd., Trade name: KBM 603) at room temperature for 20 minutes, followed by spraying nitrogen gas on the both sides of the substrate to drive off water for drying. The substrate was baked for one hour by using an oven heated to 120°C to complete treatment of the surface of the glass substrate with a silane coupling agent.

20            On the other hand, 2.7 mg of EMCS (N-(6-Maleimidocaproyloxy) succinimide: manufactured by Dojin Co., Ltd.) was weighed, and was dissolved in a solution of DMSO/ethanol (1:1) (final concentration of 0.3 mg/ml). The glass substrate subjected to the treatment with a silane coupling agent was soaked in this EMCS solution for two hours to carry out the reaction between the amino group of the silane coupling agent

covering the surface of the substrate and the succinimide group in the EMCS solution. In association with this reaction, the substrate is covered with EMCS through the silane coupling agent. In the obtained glass surface, a maleimide group derived from the EMCS exists on the surface. The glass substrate taken out after the reaction with the EMCS solution is cleaned with distilled water, and is thereafter dried with nitrogen gas. This glass substrate subjected to the surface treatment for introducing a maleimide group will be used for a binding reaction with DNA described later.

### 3. Synthesis of DNA for fixing glass substrates

Oligonucleotide having a base sequence of the following Sequence 1 is chemically synthesized for fixation on the glass substrate. This sequence 1 is a 18 mer sequence including in its central part a base sequence with a base length of 6 to code 248th and 249th amino acids in an amino acid sequence of a gene product (peptide chain) that is coded by the p 53 gene known as a tumor suppressor gene. Also, A SH group is introduced in its 5' end for fixation on the glass substrate.

Sequence 1 5' HS-GATGGGCCTCCGGTTCAT3'

The SH group is introduced by using a commercially available reagent Thiol-Modifier (manufactured by GlenResearch Co., Ltd.) on a DNA automatic synthesizing

apparatus. Subsequently, normal deprotection was carried out to recover DNA, and the DNA was purified by high speed liquid chromatography, and was then used in the following processes.

- 5           4. Discharging of DNA using a BJ printer head and binding thereof to a substrate

          The above described synthetic oligonucleotide (DNA) was dissolved in water, and the solution was diluted to the a concentration of 8  $\mu$ M using SG Clear  
10   (a solution containing 7.5% of glycerin, 7.5% of urea, 7.5% of thiodiglycol and 1% of acetylenol EH).

          100  $\mu$ l of this oligonucleotide solution was charged into the nozzle of BJ printer head BC 62 (manufactured by Canon. Inc.) with the nozzle modified  
15   so that it is suitable for a small amount of samples (discharged amount). This modified printer head was set in a plotting apparatus to perform printing over the surface of the glass substrate as an area of "solid print" of 2 mm square with the oligonucleotide  
20   solution. Furthermore, the modified printer head that was used is used for bubble jet type ink jet printing and enables printing to be performed at a resolution of 360  $\times$  720 dpi.

          Thereafter, the glass substrate coated with the  
25   oligonucleotide solution was left in a humidifier chamber for 30 minutes to carry out a reaction between the maleimide group on the surface of the substrate and

the thiol group (HS-) of oligonucleotide. Thereafter, unreacted oligonucleotide was removed. The prepared substrate to detect is a substrate with the synthetic DNA (oligonucleotide) of the above described Sequence 1  
5 bound to a predetermined section of 2 mm square on the glass substrate through covalent bond.

(Example 4)

Supply of cDNA solution to the surface of the substrate with oligonucleotide bound thereto and  
10 hybridization reaction.

From various kinds of cDNA libraries obtained from tumor tissues, p 53 gene fragments were PCR-amplified, and then only one side chains were reamplified using primers marked in advance to prepare marked single-  
15 strain cDNA for use as test samples. The hybridization reaction was carried out between this marked single strain DNA derived from the p 53 gene and the DNA probe bound on the detection substrate prepared in Example 3.

1. Preparation of test samples

20 From 64 types of cDNA libraries obtained from tumor tissues, p 53 gene fragments were obtained by the PCR reaction.

Specifically, first, all RNA samples were separated/collected from respective tissues collected with the biopsy, using Catrimox-14 (Biotechnology Co.,  
25 Ltd.). On the basis of the all RNA sample solutions, a c-DNA library was prepared using First-Strand cDNA

Synthesis Kit (manufactured by Life Science Co., Ltd.).  
A primer for amplifying p 53 genes was added to this  
cDNA library to amplify P 53 gene fragments. With this  
PCR amplification product as a template, the marked  
5 five side primer was used to carry out the PCR reaction  
(DNA synthetic reaction) to amplify only one side  
chains. By this amplification, marked single strain  
DNA derived from the p 53 gene can be prepared.

(1) Amplification of p 53 gene fragments having a  
10 T3 binding site in the terminal by the PCR method.

For using a primer for auto sequencers (Takara  
Shuzo Co., Ltd) using T3 promoters as the above  
described marked primer, a primer having a T3 site in  
the terminal and having coupled to its downstream a  
15 base sequence allowing the p 53 gene part to be  
amplified was first synthesized. The PCR reaction was  
carried out using this primer to obtain a PCR  
amplification product having a T3 promoter site coupled  
to the p 53 gene part.

20 In this example, for the five prime-end primer for  
amplifying p 53 genes, the primer with a base sequence  
having a T3 promoter site coupled to its five side (T3-  
P53) was prepared. The base sequence is shown below.

5'

25 AATTAACCCTCACTAAAGGGAACCTGAGGTGGCTCTGACTGTACCACCATCC3'

In the sequence, the underlined part on the side  
of five prime-end represents a T3 polymerase binding

site. On the other hand, for a three prime-end primer for amplification, a three prime-end primer attached in a commercially available amplification kit, "Human p 53 Amplimer Set" of CLONTECH Co., Ltd. was used. For a  
5 PCR reactive solution, "one shot LA PCR Mix" (Takara Shuzo Co., Ltd.) was used.

The solution composition in the PCR reaction has:

	one shot LA PCR Mix	25 $\mu$ l
	T3-P53 primer (20 $\mu$ M)	1 $\mu$ l
10	3' primer (20 $\mu$ M)	1 $\mu$ l
	cDNA library solution	1 $\mu$ l
	DW	22 $\mu$ l/50 $\mu$ l,

and for the PCR cycle, a condition of conducting cycles at 95°C for 30 seconds, at 55°C for 30 seconds  
15 and at 72°C for 60 seconds at 29 times after thermal denaturation at 95°C for 5 minutes and finally keeping the solution at 72°C for five minutes was used, and the reactant was stored at 4°C on a temporary basis after it was cooled.

20 After the reaction, gel electrophoresis was carried out to confirm a PCR product existing in the region of molecular weight of about 300 mer. This PCR product was purified with Micro Spin Column S200 (Pharmacia) to obtain p 53 gene fragments to which the  
25 T3 primer can be coupled (T3-linked p 53 DNA).

(2) Synthesis of marked single strain DNA using labeled T3 primers (Rho-T3).



With the p 53 gene fragment obtained in (1) as a matrix, single strain marked DNA was obtained with the PCR reaction , using a Rho-T3 primer (Takara Shuzo Co., Ltd.). The composition of the reactive solution had:

5	one shot LA PCR Mix	25 $\mu$ l
	Rho-T3primer (10 $\mu$ M)	1 $\mu$ l
	T3-linked p 53 DNA	1 $\mu$ l
	DW	23 $\mu$ l/50 $\mu$ l,

and for the reaction cycle, a condition of  
10 conducting cycles at 96°C for 30 seconds, at 50°C for  
15 seconds and at 60°C for 4 minutes 24 times was used,  
and the reactant was stored at 4°C on a temporary basis  
after it was cooled. It was purified with Micro Spin  
Column S200, and thereafter gel electrophoresis was  
15 carried out to confirm desired rhodamine labeled single  
strain DNA synthesized through the PCR reaction.

## 2. Supply of test sample solution

Sodium chloride was added in the test sample  
obtained in the above described process, namely the  
20 solution of rhodamine marked single strain DNA derived  
from the p 53 gene so that the final concentration of  
the solution was 1M. The solution of rhodamine marked  
single strain DNA derived from the p 53 gene, which had  
been prepared from 64 types of c DNA libraries was  
25 injected into each well of a 96-hole microtiter plate.  
These solutions of rhodamine marked single strain DNA  
were spotted as an arrangement of 8  $\times$  8 onto the

detection glass substrate with the DNA probe of Sequence 1 obtained in Example 3 in the form of 2 mm square, using a microarray preparing apparatus (pin system) manufactured by Cartesian Technologies. The diameter of each spot was 100  $\mu$ m.

### 3. Hybridization reaction.

This detection substrate with total 64 types of rhodamine marked single strain DNA solutions being sample specimens spotted thereon was left in a humidifier chamber set at 40°C to carry out a hybridization reaction for 3 hours. Thereafter, the detection substrate was washed with a 10 mM phosphate buffer containing 100 mM NaCl to remove test samples that had not been engaged in the formation of hybrid substances.

After the hybridization reaction, the test sample spotted in the form of a two-dimensional array of 8 x 8 was observed using an inverted fluorescence microscope equipped with a filter set for excitation light and fluorescence suitable for fluorescence marked rhodamine. For the most part of the spots, red fluorescence derived from fluorescence marked rhodamine in association with the formation of hybrid substances was observed. However, fluorescence intensity was weak for six spots and no fluorescence was observed for one spot.

For this, it can be considered that since in the p

53 gene derived from corresponding six types of tumor cells, variation occurs somewhere in the base sequence corresponding to the 248th and 249th of the amino acid sequence of the p 53 gene product (p 53 protein), the amount of formed hybrid substances is small due to its mismatch, and in association therewith, the fluorescence intensity from the fluorescence mark is weak. For the test sample in which fluorescence was not observed, it can be considered from the fact that hybrid substances were not formed that in p 53 cDNA fragments contained in the sample, deficiency occurs in the base sequence to code the above described 248th and 249th of the amino acid sequence, and consequently hybrid substances could not be formed.

(Example 5)

Preparation of array form spots of test samples on the probe matrix detection substrate with multiple oligonucleotides fixed thereon.

1. Preparation of 64 probe matrices.

Processing was performed as in the case of Example 3 to prepare a glass substrate having a maleimide group. 64 DNA of which base sequences are shown in Table 2 were printed (applied) thereon in the area of 2 mm square, respectively, using a bubble jet printer head similar to that of Example 3 to prepare a detection substrate on which sections with 64 types of prove DNAs fixed therein were arranged in a matrix

form.

Focusing attention on the 248th and 249th amino acids of the amino acid sequence of the gene product (p 53 protein) of the p 53 gene being a tumor suppressor gene, 64 DNAs of which base sequences are shown in Table 1 were selected on the basis of the base sequence to code these two amino acids so that a sequence with various kinds of base variations added thereto was obtained. Specifically, it is known that a case of frequent variation in the base sequence CGGAGG providing a base is the case where the first C of the CGG to code the 248th amino acid is changed to T, the second A is changed to G, and the third G of the AGG to code the 248th amino acid is changed to T. Thus, 64 probes were designed to provide sequences capable of being bound to base sequences with these bases at three positions varied in various kinds of forms.

Actually, it was a structure in which the total length of the probe was 18 mer, six bases including this variation were located in the center thereof, and common base sequences with base lengths of 6 were placed before and after the six bases. More specifically, the structure has a common sequence of ATGAAC from the side of the five prime-end, the base sequence of NNGAGN as a portion including the variation, and a common sequence of CCCATC on the side of three prime-end.

It was a base sequence a base sequence  
complimentary to the sequence of  
5'ATGAACNNGAGNCCCATC3'. That is, it was a probe  
expressed by 5'GATGGGNCTCNGTTCAT3'. Furthermore,  
5 since it is a DNA probe, the portion denoted by N in  
the above described base sequence refers to any one of  
A, G, C and T that are four DNA nucleic acid bases.

Table 2

1	5'-GATGGGACTCAAGTTCAT-3'	33	5'-GATGGGCCTCAAGTTCAT-3'
2	5'-GATGGGACTCAGGTTCAT-3'	34	5'-GATGGGCCTCAGGTTCAT-3'
3	5'-GATGGGACTCACGTTCAT-3'	35	5'-GATGGGCCTCACGTTCAT-3'
4	5'-GATGGGACTCATGTTCAT-3'	36	5'-GATGGGCCTCATGTTCAT-3'
5	5'-GATGGGACTCGAGTTCAT-3'	37	5'-GATGGGCCTCGAGTTCAT-3'
6	5'-GATGGGACTCGGGTTCAT-3'	38	5'-GATGGGCCTCGGGTTCAT-3'
7	5'-GATGGGACTCGCGTTCAT-3'	39	5'-GATGGGCCTCGCGTTCAT-3'
8	5'-GATGGGACTCGTGTTCAT-3'	40	5'-GATGGGCCTCGTGTTCAT-3'
9	5'-GATGGGACTCCAGTTCAT-3'	41	5'-GATGGGCCTCCAGTTCAT-3'
10	5'-GATGGGACTCCGGTTCAT-3'	42	5'-GATGGGCCTCCGGTTCAT-3'
11	5'-GATGGGACTCCC GTTTCAT-3'	43	5'-GATGGGCCTCCC GTTTCAT-3'
12	5'-GATGGGACTCCTGTTCAT-3'	44	5'-GATGGGCCTCCTGTTCAT-3'
13	5'-GATGGGACTCTAGTTCAT-3'	45	5'-GATGGGCCTCTAGTTCAT-3'
14	5'-GATGGGACTCTGGTTCAT-3'	46	5'-GATGGGCCTCTGGTTCAT-3'
15	5'-GATGGGACTCTCGTTCAT-3'	47	5'-GATGGGCCTCTCGTTCAT-3'
16	5'-GATGGGACTCTTGTTCAT-3'	48	5'-GATGGGCCTCTTGTTCAT-3'
17	5'-GATGGGGCTCAAGTTCAT-3'	49	5'-GATGGGTCTCAAGTTCAT-3'
18	5'-GATGGGGCTCAGGTTCAT-3'	50	5'-GATGGGTCTCAGGTTCAT-3'
19	5'-GATGGGGCTCACGTTCAT-3'	51	5'-GATGGGTCTCACGTTCAT-3'
20	5'-GATGGGGCTCATGTTCAT-3'	52	5'-GATGGGTCTCATGTTCAT-3'
21	5'-GATGGGGCTCGAGTTCAT-3'	53	5'-GATGGGTCTCGAGTTCAT-3'
22	5'-GATGGGGCTCGGGTTCAT-3'	54	5'-GATGGGTCTCGGGTTCAT-3'
23	5'-GATGGGGCTCGCGTTCAT-3'	55	5'-GATGGGTCTCGCGTTCAT-3'
24	5'-GATGGGGCTCGTGTTCAT-3'	56	5'-GATGGGTCTCGTGTTCAT-3'
25	5'-GATGGGGCTCCAGTTCAT-3'	57	5'-GATGGGTCTCCAGTTCAT-3'
26	5'-GATGGGGCTCCGGTTCAT-3'	58	5'-GATGGGTCTCCGGTTCAT-3'
27	5'-GATGGGGCTCCCGTTCAT-3'	59	5'-GATGGGTCTCCCGTTCAT-3'
28	5'-GATGGGGCTCCTGTTCAT-3'	60	5'-GATGGGTCTCCTGTTCAT-3'
29	5'-GATGGGGCTCTAGTTCAT-3'	61	5'-GATGGGTCTCTAGTTCAT-3'
30	5'-GATGGGGCTCTGGTTCAT-3'	62	5'-GATGGGTCTCTGGTTCAT-3'
31	5'-GATGGGGCTCTCGTTCAT-3'	63	5'-GATGGGTCTCTCGTTCAT-3'
32	5'-GATGGGGCTCTTGTTCAT-3'	64	5'-GATGGGTCTCTTGTTCAT-3'

Then, for each of the 64 types of labeled probe DNAs, a 8  $\mu$ M solution containing glycerin, urea and thiodiglycol in the final concentration of 7.5%, respectively, and acetylenol EH in the final concentration of 1% was prepared. As in the case of Example 4, using BJ Printer Head BC 62 (manufactured by Cannon Inc), different DNA probe solution was charged by 100  $\mu$ l in each of the six nozzles of the printer head, and using a plurality of such printer heads, a detection substrate with total 64 DNA probes applied to and fixed in each section of 2 mm square in the form of "solid print" and arranged in a matrix form (8  $\times$  8) was prepared. A schematic layout of the 64 DNA probes arranged in a matrix form (8  $\times$  8) on the detection substrate is shown in FIG. 7.

## 2. Preparation of array spots of test samples.

As in the case of Example 4, 64 types of labeled cDNAs were spotted in the form of the two dimensional 8  $\times$  8 array on each region of 2 mm square for fixing probes. Specifically, as schematically shown in FIG. 8, a pin system array preparing apparatus was used to form spots in the form of the two dimensional 8  $\times$  8 array on the sections arranged in a matrix form (8  $\times$  8) in which each DNA probe was fixed.

## 3. Hybridization reaction.

A hybridization reaction was carried out using conditions and procedures similar to those of Example

4. The result thereof is shown in FIG. 9. In the arrangement shown in FIG. 7, with respect to spots on probes corresponding to the base sequence of the 42nd normal gene, fluorescence intensity was weak for six spots as in the case of Example 4. Also, no fluorescence was observed for one spot. In addition thereto, it was observed that fluorescent was emitted from the spot at three points in the tenth probe region, at two points in the 41st probe region, and at one point in the 46th probe region, respectively.

Spot positions in which fluorescence in association with the formation of hybrid substances was observed in the prove region having these base sequences including variations corresponded to spot positions of weak fluorescence intensity in the probe region having the above described 42nd original base sequence. Thus, if the base sequences of the probes are compared between both the regions, the base sequence of the tenth probe is ACTCCG, the base sequence of the 41st probe is the CCTCCA, and the base sequence of the 46th probe is CCTCTG with respect to the original base sequence of CCTCCG that the 42nd probe has. For their complementary sequences, it can be understood that with respect to the CGGAGG in the 42nd probe, the CGGAGT and G were changed to T in the tenth probe, the TGGAGG and C were changed to T in the 41st probe, and the CAGAGG and G were changed to A in



the 46th probe. That is, it was confirmed that in test samples forming hybrid substances with these tenth, 41st and 46th probes, cDNA fragments contained therein derived from the p53 gene caused one base mismatch with respect to the 42nd probe due to the above described variations.

By this method, existence/not existence of variations and types thereof could be detected at the same time for all the 64 types of test samples.

10 (Example 6)

Preparation of a substrate for probe matrices partitioned by patterns.

A glass substrate with an epoxy group introduced to the surface and with black matrices for probe matrices was prepared in accordance with the following procedure.

1. Introduction of an epoxy group to the surface of the substrate.

A glass substrate made of synthetic quartz (50 mm × 50 mm) was first subjected to ultrasonic cleaning using a 2% sodium hydrate solution, and was then subjected to UV ozone processing to clean the surface. A 50% methanol solution containing 1% of silane coupling agent (trade name: KBM 403; manufactured by The Shin-Etsu Chemical Co., Ltd.) containing a silane compound having an epoxy group bonded thereto ( $\gamma$ -glycidoxypopyltrimethoxysilane) was stirred at room

temperature for three hours to perform preliminary treatment for hydrolyzing the methoxy group in the silane compound. This solution already subjected to the hydrolysis treatment was applied to the surface of the above described clean substrate with a spin coater, and was heated and dried at 100°C for 5 minutes to form a binding coating of the silane coupling agent on the surface of the substrate. Through the formation of this coating, the epoxy group contained in the silane compound was introduced to the surface of the substrate.

## 2. Formation of black matrices.

Then, A DEEP-UV resist containing carbon black (negative type resist for black matrices) (trade name: BK-739P; manufactured by Nippon Steel Chemical Co., Ltd.) was applied on the surface of the substrate with a spin coater so that the film thickness after curing was 5 µm, and it was heated for curing on a hotplate at 80°C for 5 minutes. By proximity exposure using a DEEP-UV aligner, a pattern was exposed to light using as an exposure mask a mask for negatives with patterning applied to a region of 10 mm × 10 mm so that the distance X between adjacent wells was 100 µm and the outer shape of the well was a square of 1 mm × 1 mm. Then, development was carried out with a developer of inorganic aqueous alkaline solution using a spin drier, and the substrate was washed with pure water to

remove the developer completely. Then, it was briefly dried using the spin drier, and was thereafter heated in a clean oven at 180°C for 30 minutes to fully cure the resist. As a whole, a substrate with 400 wells arranged in a predetermined arrangement and black matrices (resist walls) partitioning adjacent wells was obtained. Furthermore, the internal volume of each well is calculated as 5  $\mu$ l if the thickness of solution is 5  $\mu$ m. Also, in the surface of the prepared black matrix, the angle of contact to water was 93 degrees and wettability with water was significantly low, while in the bottom of the well, the angle of contact to water was 35 degrees and the wettability with water was high.

### 3. Fixation of probe DNA.

64 oligonucleotides of 18 mer with an amino group bound to the hydroxyl group of the five prime-end through a phosphate group and hexamethylene were prepared as DNA probes. The 64 probes are same as those prepared in Example 5 as to base sequences, but are different in the sense that an amino group is introduced in its five prime-end instead of a thiol group.

5  $\mu$ l of solution of these DNA probes was injected into each well under a microscope, and was left in a humidified chamber to allow the probe bind to the substrate through the reaction between the amino group

of the five primer-end and the epoxy group on the substrate.

(Example 7)

Analysis of cDNA derived from the p 53 gene that  
5 has been prepared from mRNA, using the probe matrix  
substrate partitioned by the pattern that has been  
prepared in Example 6.

As in the case of Example 4, 64 types of labeled  
cDNAs were spotted in each probe region of 2 mm square  
10 as an arrangement of 8 x 8 spots as shown in FIG. 8,  
using a pin system array preparing apparatus.

A hybridization reaction was carried out by a  
method similar to that of example 4.

The obtained result was similar to that of Example  
15 5.